




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 239/95, 401/04, A61K 31/505	A1	(11) International Publication Number: WO 97/20820 (43) International Publication Date: 12 June 1997 (12.06.97)
(21) International Application Number: PCT/EP96/05055 (22) International Filing Date: 18 November 1996 (18.11.96) (30) Priority Data: 08/566,349 1 December 1995 (01.12.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/566,349 (CIP) Filed on 1 December 1995 (01.12.95) (71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basle (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): RÜEGER, Heinrich [CH/CH]; Alemannenweg 6, CH-4112 Flüh (CH). SCHMIDLIN, Tibur [CH/CH]; Friedensgasse 36, CH-4056 Basle (CH). RIGOLLIER, Pascal [FR/FR]; 2, rue Sainte-Catherine, F-68100 Mulhouse (FR). YAMAGUCHI, Yasuchika [JP/CH]; Tellstrasse 44/2, CH-4053 Basle (CH). TINTELNOT-BLOMLEY, Marina [DE/DE]; Röttlerstrasse 1, D-79689 Maulburg (DE). SCHILLING, Walter [CH/CH];		Im Muspenacker, CH-4204 Himmelried (CH). CRISCIONE, Leoluca [IT/CH]; Kirchstrasse 15, CH-4313 Möhlin (CH). (74) Common Representative: NOVARTIS AG; Patent and Trade-mark Dept., Klybeckstrasse 141, CH-4002 Basle (CH). (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: HETEROARYL COMPOUNDS**(57) Abstract**

The invention relates to a method of treatment of disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal, including man, in need of such treatment a therapeutically effective amount of a compound of formula (I) in which the variables are as defined and relates to new compounds of formula (I) or a salt thereof, to pharmaceutical compositions, and to the manufacture of new compounds of formula (I) and salts thereof.



(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Heteroaryl Compounds

Background of the Invention


Neuropeptide Y (NPY) is a member of the pancreatic polypeptide family of peptides and is one of the most abundant and widely distributed peptides at the central and peripheral nervous system. NPY acts as a neurotransmitter playing an important role in the regulation of various diseases. Intensive evaluations lead to the finding that multiple NPY receptors are existing being responsible for different physiological and pharmacological activities. Recently, a new NPY receptor subtype has been characterized and cloned, designated as Y5 receptor. It has been demonstrated that the pharmacological function associated with Y5 relates, for example, to obesity and eating disorders. Accordingly, the provision of compounds which act as antagonists of this receptor subtype represents a promisable approach in the regulation of diseases or disorders, such as obesity and eating/food intake disorders.

Summary of the Invention

The invention relates to a method of treatment of disorders and diseases associated with NPY receptor subtype Y5, to pharmaceutical compositions and to new compounds having Y5 antagonistic properties.

Detailed Description of the Invention

The invention relates to a method of treatment and prophylaxis of disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal, including man, in need of such treatment a therapeutically effective amount of a compound of formula (I)



in which

- 2 -

alk represents a single bond or lower alkylene;

the integer n is 0 or 1;

R₁ represents

- (i) hydrogen, halogen, nitro, cyano, lower alkyl, lower alkenyl, lower alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, by lower alkoxy, by amino, by substituted amino, by carboxy, by lower alkoxycarbonyl, by (carbocyclic or heterocyclic) aryl-lower alkoxycarbonyl, by carbamoyl, or by N-substituted carbamoyl;
- (ii) amino or substituted amino;
- (iii) hydroxy, lower alkoxy, lower alkenyloxy, lower alkynyloxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, C₃-C₈-cycloalkoxy, C₃-C₈-cycloalkyl-lower alkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, lower alkoxycarbonyl-oxy, (carbocyclic or heterocyclic) aryl-lower alkoxycarbonyl-oxy, aminocarbonyl-oxy, or N-substituted aminocarbonyl-oxy;
- (iv) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (v) carbamoyl or N-substituted carbamoyl;
- (vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-O-R, -NR₀₁-CO-R, -NR₀₁-CO-NR₀₁-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R and R₀₁ being as defined below, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O, S(O)_n or NR₀ or which may be condensed at two adjacent carbon atoms with a benzene ring}]; or
- (vii) an element of formula -X₁(X₂)(X₃) wherein, (a) if X₁ is -CH-, X₂ together with X₃ represent a structural element of formula -X₄-(CO)_p-(CH₂)_o-, -(CH₂)_q-X₄-(CO)_p-(CH₂)_r-, or -(CH₂)_s-X₄-CO-(CH₂)_t-; or, (b) if X₁ is -N-, X₂ together with X₃ represent a structural element of formula -CO-(CH₂)_u-; [X₄ being -CH₂-, -N(R₀₁)- or -O-; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X₄ is different from -CH₂-];

R₂ and R₃, independently of one another, represent

- (i) hydrogen, lower alkyl, lower alkenyl, lower alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl,

(carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, N-substituted carbamoyl, and $-S(O)_n-R$;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O, $S(O)_n$, NR_0 or which may be condensed at two adjacent carbon atoms with a benzene ring];

X represents N or CH;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, lower alkenyl, lower alkynyl, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, lower alkynyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, (carbocyclic or heterocyclic) aroyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iv) amino, substituted amino;

(v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C_3 - C_8 -cycloalkyl, by C_3 - C_8 -cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, $S(O)_n$ or NR_0 or which may be condensed at two adjacent carbon atoms with a benzene

- 4 -

ring], or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by $-\text{CO}-(\text{O})_v-\text{R}$ and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R_0 represents hydrogen, lower alkyl, lower alkenyl, lower alkinyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower alkanoyl, (carbocyclic or heterocyclic) aroyl, $-\text{SO}_2-\text{R}$, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;

wherein in each case R_{01} represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy; or a pharmaceutically acceptable salt thereof; and relates to new compounds of formula (I) or a salt thereof, to pharmaceutical compositions, and to the manufacture of new compounds of formula (I) and salts thereof.

The compounds (I) can be present as salts, in particular pharmaceutically acceptable salts. If the compounds (I) have, for example, at least one basic centre, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, a phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as C_1-C_4 -alkanecarboxylic acids which are unsubstituted or substituted, for example, by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, for example aspartic or glutamic acid, or such as benzoic acid, or with organic sulfonic acids, such as C_1-C_4 -alkane- or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methane- or p-toluenesulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic centre. The compounds (I) having at least one acid group (for example COOH) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine,

a mono-, di- or tri-lower alkylamine, for example ethyl-, tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethylpropylamine, or a mono-, di- or trihydroxy lower alkylamine, for example mono-, di- or triethanolamine. Corresponding internal salts may furthermore be formed, if a compound of formula comprises e.g. both a carboxy and an amino group. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds (I) or their pharmaceutically acceptable salts, are also included.

(Carbocyclic or heterocyclic) aryl in (carbocyclic or heterocyclic) aryl or aryloxy, respectively, represents, for example, phenyl, biphenyl, naphthyl or an appropriate 5- or 6-membered and monocyclic radical or an appropriate bicyclic heteroaryl radical which, in each case, have up to four identical or different hetero atoms, such as nitrogen, oxygen or sulfur atoms, preferably one, two, three or four nitrogen atoms, an oxygen atom or a sulfur atom. Appropriate 5-membered heteroaryl radicals are, for example, monoaza-, diaza-, triaza-, tetraaza-, monooxa- or monothia-cyclic aryl radicals, such as pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furyl and thienyl, while suitable appropriate 6-membered radicals are in particular pyridyl. Appropriate bicyclic heterocyclic aryls are, for example, indolyl, indazolyl, benzofuryl, benzothiophenyl, benzimidazolyl, quinoliny, isoquinoliny, or quinazoliny. Appropriate aromatic radicals, including ring A, are radicals which may be monosubstituted or polysubstituted, for example di- or trisubstituted, for example by identical or different radicals, for example selected from the group as given above. Preferred substituents of corresponding aryl radicals (including of ring A) are, for example, halogen, lower alkyl, halo-lower alkyl, lower alkoxy, oxy-lower alkylene-oxy, hydroxy, hydroxy-lower alkoxy, and lower alkoxy-lower alkoxy.

(Carbocyclic or heterocyclic) aroyl is in particular benzoyl, naphthoyl, furoyl, thenoyl, or pyridoyl.

(Carbocyclic or heterocyclic) aryl-lower alkanoyl in (carbocyclic or heterocyclic) aryl-lower alkanoyloxy or (carbocyclic or heterocyclic) aryl-lower alkanoyl is in particular phenyl-lower alkanoyl, naphthyl-lower alkanoyl, or pyridyl-lower alkanoyl.

(Carbocyclic or heterocyclic) aryl-lower alkyl is in particular phenyl-, naphthyl- or pyridyl-lower alkyl.

(Carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl is in particular phenyl-, naphthyl- or pyridyl-lower alkoxy.

Lower alkyl which substituted by halogen, hydroxy, lower alkoxy, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, or amino is in particular halo-lower alkyl, hydroxy-lower alkyl, lower alkoxy-lower alkyl, phenyloxy-, naphthyloxy- or pyridyloxy-lower alkyl, phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl, amino-lower alkyl, or N- or N,N- substituted amino-lower alkyl.

An amino group which is mono-substituted by lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl is in particular lower alkyl-amino, C₃-C₈-cycloalkyl-amino, C₃-C₈-cycloalkyl-loweralkyl-amino, phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-amino, phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl-amino.

An amino group which is, independently of one another, di-substituted by lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, or (carbocyclic or heterocyclic) aryl-lower alkyl is in particular di-lower alkyl-amino, di-C₃-C₈-cycloalkyl-amino, di-(C₃-C₈-cycloalkyl-lower alkyl)-amino, di-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-amino, di-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl)-amino, lower alkyl-C₃-C₈-cycloalkyl-amino, lower alkyl-(C₃-C₈-cycloalkyl-lower alkyl)-amino, lower alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-amino, lower alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl)-amino.

Lower alkyl which is substituted by carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, carbamoyl in which the amino group is mono-substituted or, independently of one another, di-substituted by lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, and carbamoyl in which the amino group is di-substituted by lower alkylene [which may be interrupted by O, S(O)_n, NR₀, the integer n being 0, 1 or 2 and R₀ being hydrogen, lower alkyl, (carbocyclic or

heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower alkanoyl, (carbocyclic or heterocyclic) aroyl, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{-R}$ and R being lower alkyl, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryl, or (carbocyclic or heterocyclic) aryl-lower alkyl] is in particular carboxy-lower alkyl, lower alkoxy-carbonyl-lower alkyl, lower alkoxy-lower alkoxy-carbonyl-lower alkyl, (phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-lower alkoxy-carbonyl-lower alkyl, carbamoyl-lower alkyl, or N- or N,N-substituted carbamoyl-lower alkyl.

Lower alkoxy which substituted by halogen, hydroxy, lower alkoxy, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, or amino is in particular halo-lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, phenyloxy-, naphthyloxy- or pyridyloxy-lower alkyl, phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkoxy, amino-lower alkoxy, or corresponding N- or N,N- substituted amino-lower alkoxy.

Lower alkoxy which is substituted by carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, carbamoyl in which the amino group is mono-substituted or, independently of one another, di-substituted by lower alkyl, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, and carbamoyl in which the amino group is di-substituted by lower alkylene [which may be interrupted by O, $\text{S}(\text{O})_n$, NR_0 , the integer n being 0, 1 or 2 and R_0 being hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower alkanoyl, (carbocyclic or heterocyclic) aroyl, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{-R}$ and R being lower alkyl, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryl, or (carbocyclic or heterocyclic) aryl-lower alkyl] is in particular carboxy-lower alkoxy, lower alkoxy-carbonyl-lower alkoxy, lower alkoxy-lower alkoxy-carbonyl-lower alkoxy, (phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-lower alkoxy-carbonyl-lower alkoxy, carbamoyl-lower alkoxy, or corresponding N- or N,N-substituted carbamoyl-lower alkoxy.

Substituted lower alkyl or lower alkoxy, respectively, is mono- or poly-substituted, e.g. di- or tri-substituted.

The group of formula $-N(R_2)(R_3)$ in which R_2 and R_3 together represent lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring represents, for example, lower alkylene-phenylene-lower alkylene-amino, such as 3,4-dihydro-1*H*-isoquinolin-2-yl.

The general definitions used above and below, unless defined differently, have the following meanings:

The expression "lower" means that corresponding groups and compounds, in each case, in particular comprise not more than 7, preferably not more than 4, carbon atoms.

Halogen is in particular halogen of atomic number not more than 35, such as fluorine, chlorine or bromine, and also includes iodine.

Lower alkyl is in particular C_1 - C_7 -alkyl, for example methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, sec-butyl, tert-butyl, and also includes corresponding pentyl, hexyl and heptyl radicals. C_1 - C_4 -alkyl is preferred.

Lower alkenyl is in particular C_3 - C_7 -alkenyl and is, for example, 2-propenyl or 1-, 2- or 3-butenyl. C_3 - C_5 -alkenyl is preferred.

Lower alkynyl is in particular C_3 - C_7 -alkynyl and is preferably propargyl.

Lower alkoxy is in particular C_1 - C_7 -alkoxy and is, for example, methoxy, ethoxy, *n*-propyloxy, isopropyloxy, *n*-butyloxy, isobutyloxy, sec-butyloxy, tert-butyloxy and also includes corresponding pentyloxy, hexyloxy and heptyloxy radicals. C_1 - C_4 -alkoxy is preferred.

Lower alkenyloxy is in particular C_3 - C_7 -alkenyloxy, preferably allyloxycarbonyl, while lower alkynyloxy is in particular C_3 - C_5 -alkynyloxy, such as propargyloxy.

Oxy-lower alkylene-oxy is in particular oxy- C_1 - C_4 -alkylene-oxy, preferably oxy-methylene-oxy or oxy-ethylene-oxy.

Lower alkanoyloxy is in particular C₂-C₇-alkanoyloxy, such as acetyloxy, propionyloxy, butyryloxy, isobutyryloxy or pivaloyloxy. C₂-C₅-alkanoyloxy is preferred.

Lower alkanoyl is in particular C₂-C₇-alkanoyl, such as acetyl, propionyl, butyryl, isobutyryl or pivaloyl. C₂-C₅-alkanoyl is preferred.

Naphthoyl is 1- or 2-naphthoyl, furoyl 2- or 3-furoyl, thenoyl 2- or 3-thenyl, and pyridoyl 2-, 3-, or 4-pyridoyl.

C₃-C₈-Cycloalkyl is, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Cyclopentyl and cyclohexyl are preferred.

C₃-C₈-Cycloalkyl-lower alkyl is in particular C₃-C₈-cycloalkyl-C₁-C₄-alkyl, in particular C₃-C₆-cycloalkyl-C₁-C₂-alkyl. Preferred is cyclopropylmethyl, cyclopentylmethyl or cyclohexylmethyl.

C₃-C₈-Cycloalkoxy is, for example, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and cycloheptyloxy. Cyclopentyloxy and cyclohexyloxy are preferred.

C₃-C₈-Cycloalkyl-lower alkoxy is in particular C₃-C₈-cycloalkyl-C₁-C₄-alkoxy, in particular C₃-C₆-cycloalkyl-C₁-C₂-alkoxy. Preferred is cyclopropylmethoxy, cyclopentylmethoxy or cyclohexylmethoxy.

Lower alkylene is in particular C₁-C₇-alkylene, in particular C₁-C₅-alkylene, and is straight-chain or branched and is in particular methylene, ethylene, propylene and butylene and also 1,2-propylene, 2-methyl-1,3-propylene, 3-methyl-1,5-pentylene and 2,2-dimethyl-1,3-propylene. C₃-C₅-alkylene is preferred. In case of alk₁ or alk₂, respectively, lower alkylene preferably is -(CH₂)_p- the integer p being 1-3. Lower alkylene in an substituted amino group preferably is 1,2-ethylene, 1,3-propylene, 1,4-butylene, 1,5-pentylene, 1,6-hexylene, 2-methyl-1,3-propylene, or 2-methyl-butylene, or 3-methyl-1,5-pentylene.

Amino which di-substituted by lower alkylene is in particular C₃-C₇-alkyleneamino, preferably 1-azidino, 1-pyrrolidino or 1-piperidino.

Amino which di-substituted by lower alkylene which is interrupted by O, S(O)_n or NR₀ is in particular morpholino, thiomorpholino or the mono- or di-oxide thereof, or 4-R₀-piperazino.

Lower alkanesulfonyl is in particular C₁-C₄-alkoxy-C₁-C₅-alkoxycarbonyl, preferably ethoxyethoxycarbonyl, methoxyethoxycarbonyl and isopropoxyethoxycarbonyl.

Lower alkoxycarbonyl is in particular C₂-C₈-alkoxycarbonyl and is, for example, methoxy-, ethoxy-, propyloxy- or pivaloyloxy-carbonyl. C₂-C₅-alkoxycarbonyl is preferred.

Lower alkoxy-lower alkoxy-carbonyl is in particular C₁-C₄-alkoxy-C₂-C₅-alkoxycarbonyl and is, for example, methoxy- or ethoxy-ethoxy-alkoxycarbonyl.

Hydroxy-lower alkyl is in particular hydroxy-C₁-C₄-alkyl, such as hydroxymethyl, 2-hydroxyethyl or 3-hydroxypropyl.

Hydroxy-lower alkoxy is in particular hydroxy-C₁-C₄-alkoxy, such as hydroxymethyl, 2-hydroxyethyl or 3-hydroxypropyl. Furthermore, hydroxy-lower alkyl may exhibit two hydroxy groups, such as 3-hydroxy-1-hydroxymethyl-propyl.

Lower alkoxy-lower alkoxy is in particular C₁-C₄-alkoxy-C₁-C₄-alkoxy and is, for example, 2-methoxyethoxy, 2-ethoxyethoxy, 2-n-propyloxyethoxy or ethoxymethoxy.

Amino which di-substituted by lower alkylene and is condensed at two adjacent carbon atoms with a benzene ring is in particular C₂-C₆-cycloalkylenemino which is condensed at two adjacent carbon atoms with a benzene ring. Preferred is indolin-1-yl or 1,2,3,4-tetrahydro-quinolin-1-yl.

Halo-lower alkyl is in particular halo-C₁-C₄-alkyl, such as trifluoromethyl, 1,1,2-trifluoro-2-chloroethyl or chloromethyl.

Halo-lower alkoxy is in particular halo-C₁-C₄-alkoxy, such as trifluoromethoxy, 1,1,2-trifluoro-2-chloroethoxy or chloromethoxy.

Phenyloxy-, naphthyloxy- or pyridyloxy-lower alkyl is in particular phenyloxy-, naphthyloxy- or pyridyloxy-C₁-C₄-alkyl, such as phenoxy-methyl, 2-phenoxy-ethyl, 1- or 2-naphthyloxy-methyl, or 2-, 3-, or 4-pyridyloxy-methyl.

Phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl is in particular phenyl-, naphthyl- or pyridyl-C₁-C₄-alkyl, such as phenyl-methyl, 2-phenyl-ethyl, 1- or 2-naphthyl-methyl, or 2-, 3-, or 4-pyridyl-methyl.

Phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkoxy is in particular phenyl-, naphthyl- or pyridyl-C₁-C₄-alkoxy, such as phenyl-methoxy, 2-phenyl-ethoxy, 1- or 2-naphthyl-methoxy, or 2-, 3-, or 4-pyridyl-methoxy.

Naphthyl is in particular 1- or 2-naphthyl; furyl 2- or 3-furyl; thienyl 2- or 3-thienyl; pyridyl 2-, 3- or 4-pyridyl, indolyl, indazolyl e.g. 6-1(H)-indazolyl, benzofuryl e.g. 2-, 3- or 5-benzofuranyl, benzothienyl e.g. 2-, 3-, or 5-benzothienyl, benzimidazolyl e.g. 1-, 2- or 5-benzimidazolyl, quinolinyln e.g. 2-, 4-, 5-, 6-, 7-, or 8-quinolinyln, isoquinolinyln e.g. 1-, 3-, 4-, or 6-isoquinolinyln, or quinazolinyln e.g. 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyln.

Amino-lower alkyl is in particular amino-C₁-C₇-alkyl, preferably amino-C₁-C₄-alkyl, such as aminomethyl, 2-aminoethyl or 3-aminopropyl.

Lower alkylamino is in particular C₁-C₇-alkylamino and is, for example, methyl-, ethyl-, n-propyl- and isopropyl-amino. C₁-C₄-alkylamino is preferred.

C₃-C₈-Cycloalkyl-amino is in particular C₃-C₆-cycloalkyl-amino and is, for example, cyclopropyl-, cyclopentyl and cyclohexyl-amino.

C₃-C₈-Cycloalkyl-lower alkylamino is in particular C₃-C₈-cycloalkyl-C₁-C₇-alkylamino and is, for example, cyclopropylmethyl-amino or cyclohexylmethyl-amino. C₃-C₈-Cycloalkyl-C₁-C₄-alkylamino is preferred.

Phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl-amino is in particular phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-C₁-C₄-alkyl-amino, preferably benzyl-amino, 2-phenethyl-amino, 1- or 2-naphthylmethyl-amino, or 2-, 3-, or 4-pyridylmethyl-amino.

Di-lower alkylamino is in particular di-C₁-C₄-alkylamino, such as dimethyl-, diethyl-, di-n-propyl-, methylpropyl-, methylethyl-, methylbutyl-amino and dibutylamino.

Di-C₃-C₈-cycloalkyl-amino is in particular di-C₃-C₆-cycloalkylamino, preferably cyclopropylamino, cyclopentylamino or cyclohexylamino.

Di-(C₃-C₈-cycloalkyl-lower alkyl)-amino is in particular di-(C₃-C₆-cycloalkyl-C₁-C₄-alkyl)-amino, preferably cyclopropylmethyl-amino, cyclopentylmethyl-amino or cyclohexylmethyl-amino.

Di-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl)-amino is in particular di-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-C₁-C₄-alkyl)-amino, preferably di-benzyl-amino, di-(2-phenethyl)-amino, di-(1- or 2-naphthylmethyl)-amino, or di-(2-, 3-, or 4-pyridylmethyl)-amino.

Lower alkyl-C₃-C₈-cycloalkyl-amino is in particular C₁-C₄-alkyl-C₃-C₆-cycloalkyl-amino, preferably methyl-cyclopropyl-amino, methyl-cyclopentyl-amino or methyl-cyclohexyl-amino.

Lower alkyl-(C₃-C₈-cycloalkyl-lower alkyl)-amino is in particular C₁-C₄-alkyl-(C₃-C₆-cycloalkyl-C₁-C₄-alkyl)amino, preferably methyl-cyclopropylmethyl-amino, methyl-cyclopentylmethyl-amino or methyl-cyclohexylmethyl-amino.

Lower alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-amino is in particular C₁-C₄-alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-amino, such as (m)ethyl-phenyl-amino.

Lower alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-lower alkyl)-amino is in particular C₁-C₄-alkyl-(phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl-C₁-C₄-alkyl)-amino, such as (m)ethyl-benzyl-amino or (m)ethyl-(2-phenethyl)-amino.

Carboxy-lower alkyl is in particular carboxy-C₁-C₄-alkyl, such as carboxy-methyl, 2-carboxy-ethyl, or 3-carboxy-propyl.

Lower alkoxy-carbonyl-lower alkyl is in particular C₂-C₅-alkoxycarbonyl-C₁-C₄-alkyl, such as (m)ethoxycarbonyl-methyl, 2-(m)ethoxycarbonyl-ethyl or 2-pivaloyl-ethyl.

Lower alkoxy-lower alkoxy-carbonyl-lower alkyl is in particular C₁-C₄-alkoxy-C₂-C₅-alkoxycarbonyl-C₁-C₄-alkyl, such as 2-methoxy-ethoxycarbonyl-methyl or 2-(2-ethoxy-ethoxycarbonyl)-ethyl.

(Phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-lower alkoxycarbonyl-lower alkyl is in particular (phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-C₂-C₅-alkoxycarbonyl-C₁-C₄-alkyl, such as benzyloxycarbonyl-methyl or 2-(2-phenethyloxy-carbonyl)-ethyl.

Carbamoyl-lower alkyl is in particular carbamoyl-C₁-C₄-alkyl, such as carbamoyl-methyl, 2-carbamoyl-ethyl or 3-carbamoyl-propyl.

Amino-lower alkoxy is in particular amino-C₁-C₄-alkoxy, such as aminomethoxy, 2-aminoethoxy, or 3-amino-propoxy.

Carboxy-lower alkoxy is in particular carboxy-C₁-C₄-alkoxy, such as carboxy-methoxy, 2-carboxy-ethoxy, or 3-carboxy-propyloxy.

Lower alkoxy-carbonyl-lower alkoxy is in particular C₂-C₅-alkoxycarbonyl-C₁-C₄-alkoxy, such as (m)ethoxycarbonyl-methoxy, 2-methoxycarbonyl-ethyl, or 2-(2-ethoxycarbonyl)-ethyl.

Lower alkoxy-lower alkoxy-carbonyl-lower alkoxy is in particular C₁-C₄-alkoxy-C₂-C₅-alkoxycarbonyl-C₁-C₄-alkoxy, such as (m)ethoxymethoxycarbonyl-methoxy, 2-ethoxy-methoxycarbonyl-ethyl, or 2-[(2-ethoxy-ethoxycarbonyl)]-ethyl.

(Phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-lower alkoxycarbonyl-lower alkoxy is in particular (phenyl-, naphthyl-, furyl-, thienyl-, or pyridyl)-C₂-C₅-alkoxycarbonyl-C₁-C₄-alkoxy,

such as benzyloxycarbonyl-methoxy, phenethyloxycarbonyl-methoxy, 2-(benzyloxycarbonyl)-ethoxy, or 2-(2-phenethyloxycarbonyl)-ethoxy.

Carbamoyl-lower alkoxy is in particular carbamoyl-C₁-C₄-alkoxy, such as carbamoyl-methoxy, 2-carbamoyl-ethoxy, or 3-carbamoyl-propyloxy.

Obesity, for example, is a wide-spread phenomena which e.g. causes a variety of pathological symptoms or influences the overall state of health. Also associated therewith are considerable socio-economic investments and a heavy financial burden for managed health care organisations. The problem to be solved is to present an approach to systemically treat obesity or related diseases or disorders. Surprisingly, it has been manifested that the modulation of the NPY receptor subtype Y5 leads to a control of the eating behavior.

Extensive pharmacological investigations have shown that the compounds (I) and their pharmaceutically acceptable salts, for example, are useful as antagonists of the neuropeptide Y5 receptor subtype.

Neuropeptide Y (NPY) is a member of the pancreatic polypeptide family with wide-spread distribution throughout the mammalian nervous system. NPY and its relatives (peptide YY or PYY, and pancreatic polypeptide or PP) elicit a broad range of physiological effects through activation of at least five G protein-coupled receptor subtypes known as Y1, Y2, Y3, Y4 (or PP), and the "atypical Y1". The role of NPY as the most powerful stimulant of feeding behavior yet described is thought to occur primarily through activation of the hypothalamic "atypical Y1" receptor. This receptor is unique in that its classification is based solely on feeding behavior data, rather than radioligand binding data, unlike the Y1, Y2, Y3, and Y4 (or PP) receptors, each of which are described previously in both radioligand binding and functional assays. ¹²⁵I-PYY-based expression cloning technique may be used to isolate a rat hypothalamic cDNA encoding an "atypical Y1" receptor referred to herein as the Y5 subtype. Y5 homolog may be isolated and characterized of from human hippocampus. Protein sequence analysis reveals that the Y5 receptor belongs to the G protein-coupled receptor superfamily. Both the human and rat homolog display $\leq 42\%$ identity in transmembrane domains with the previously cloned "Y-type" receptors. Rat brain localization studies using in situ hybridization techniques verify the existence of Y5 receptor mRNA in rat hypothalamus. Pharmacological evaluation reveals the

following similarities between the Y5 and the "atypical Y1" receptor. 1) Peptides bind to the Y5 receptor with a rank order of potency identical to that described for the feeding response: $\text{NPY}^3 \text{ NPY}_{2-36} = \text{PYY} = [\text{Leu}^{31}, \text{Pro}^{34}]\text{NPY} \gg \text{NPY}_{13-36}$. 2) The Y5 receptor is negatively coupled to cAMP accumulation, as has been proposed for the "atypical Y1" receptor. 3) Peptides activate the Y5 receptor with a rank order of potency identical to that described for the feeding response. 4) The reported feeding "modulator" $[\text{D-Trp}^{32}]\text{NPY}$ binds selectively to the Y5 receptor and subsequently activated the receptor. 5) Both the Y5 and the "atypical Y1" receptors are sensitive to deletions or modifications in the midregion of NPY and related peptide ligands.

The peptide neurotransmitter neuropeptide Y (NPY) is a 36 amino acid member of the pancreatic polypeptide family with widespread distribution throughout the mammalian nervous system. NPY is considered to be the most powerful stimulant of feeding behavior yet described (Clark, J.T., Kalra, P.S., Crowley, W.R., and Kalra, S.P. (1984). Neuropeptide Y and human pancreatic polypeptide stimulate feeding behavior in rats. Endocrinology 115: 427-429, 1984; Levine, A.S., and Morley, J.E. (1984). Neuropeptide Y: A potent inducer of consummatory behavior in rats. Peptides 5: 1025-1029; Stanley, B.G., and Leibowitz, S.F.; (1984) Neuropeptide Y: Stimulation of feeding and drinking by injection into the paraventricular nucleus. Life Sci. 35: 2635-2642). Direct injection into the hypothalamus of satiated rats, for example, can increase food intake up to 10-fold over a 4-hour period (Stanley, B.G., Magdalin, W., Seirafi, A., Nguyen, M.M., and Leibowitz, S.F. (1992). Evidence for neuropeptide Y mediation of eating produced by food deprivation and for a variant of the Y_1 receptor mediating this peptide's effect. Peptides 13: 581-587). The role of NPY in normal and abnormal eating behavior, and the ability to interfere with NPY-dependent pathways as a means to appetite and weight control, are areas of great interest in pharmacological and pharmaceutical research (Sahu and Kalra, 1993; Dryden, S., Frankish, H., Wang, Q., and Williams, G. (1994). Neuropeptide Y and energy balance: one way ahead for the treatment of obesity? Eur. J. Clin. Invest. 24: 293-308). Any credible means of studying or controlling NPY-dependent feeding behavior, however, must necessarily be highly specific as NPY can act through at least 5 pharmacologically defined receptor subtypes to elicit a wide variety of physiological functions (Dumont, Y., J.-C. Martel, A. Fournier, S. St-Pierre, and R. Quirion. (1992). Neuropeptide Y and neuropeptide Y receptor subtypes in brain and peripheral tissues. Progress in Neurobiology 38: 125-167). It is therefore vital that knowledge of the molecular biology and structural diversity of the individual receptor subtypes be understood as part of a rational drug design approach to develop subtype selective

compounds. A brief review of NPY receptor pharmacology is summarized below and also in Table 1.

TABLE 1: Pharmacologically defined receptors for NPY and related pancreatic polypeptides.

Rank orders of affinity for key peptides (NPY, PYY, PP, [Leu³¹,Pro³⁴]NPY, NPY₂₋₃₆, and NPY₁₃₋₃₆) are based on previously reported binding and functional data (Schwartz, T.W., J. Fuhlendorff, L.L.Kjems, M.S. Kristensen, M. Vervelde, M. O'Hare, J.L. Krstenansky, and B. Bjornholm. (1990). Signal epitopes in the three-dimensional structure of neuropeptide Y. Ann. N.Y. Acad. Sci. 611: 35-47; Wahlestedt, C., Karoum, F., Jaskiw, G., Wyatt, R.J., Larhammar, D., Ekman, R., and Reis, D.J. (1991). Cocaine-induced reduction of brain neuropeptide Y synthesis dependent on medial prefrontal cortex. Proc. Natl. Acad. Sci. 88: 2978-2082; Dumont, Y., J.-C. Martel, A. Fournier, S. St-Pierre, and R. Quirion. (1992). Neuropeptide Y and neuropeptide Y receptor subtypes in brain and peripheral tissues. Progress in Neurobiology 38: 125-167; Wahlestedt, C., and D.J. Reis. (1993). Neuropeptide Y-Related Peptides and Their Receptors--Are the Receptors Potential Therapeutic Targets? Ann. Rev. Pharmacol. Tox. 32: 309-352). Missing peptides in the series reflect a lack of published information.

TABLE 1

Receptor	Affinity (pK _i or pEC ₅₀)					
	11 to 10	10 to 9	9 to 8	8 to 7	7 to 6	< 6
Y1	NPY PYY [Leu ³¹ ,Pro ³⁴] NPY		NPY ₂₋₃₆	NPY ₁₃₋₃₆	PP	
Y2		PYY NPY NPY ₂₋₃₆	NPY ₁₃₋₃₆			[Leu ³¹ ,Pro ³⁴] NPY PP
Y3		NPY	[Pro ³⁴] NPY	NPY ₁₃₋₃₆ PP		PYY
Y4	PP	PYY [Leu ³¹ ,Pro ³⁴] NPY	NPY NPY ₂₋₃₆	NPY ₁₃₋₃₆		
Y5		PYY NPY NPY ₂₋₃₆ [Leu ³¹ ,Pro ³⁴] NPY		NPY ₁₃₋₃₆		

NPY Receptor Pharmacology

NPY receptor pharmacology has historically been based on structure/activity relationships within the pancreatic polypeptide family. The entire family includes the namesake pancreatic polypeptide (PP), synthesized primarily by endocrine cells in the pancreas; peptide YY (PYY), synthesized primarily by endocrine cells in the gut; and NPY, synthesized primarily in neurons (Michel, M.C. (1991). Receptors for neuropeptide Y: multiple subtypes and multiple second messengers. Trends Pharmacol.: 12: 389-394; Dumont et al., 1992; Wahlestedt and Reis, 1993). All pancreatic polypeptide family members share a compact structure involving a "PP-fold" and a conserved C-terminal hexapeptide ending in Tyr³⁶ (or Y³⁶ in the single letter code). The striking conservation of Y³⁶ has prompted the reference to the pancreatic polypeptides' receptors as "Y-type" receptors (Wahlestedt, C., L. Edvinsson, E. Ekblad, and R. Hakanson. Effects of neuropeptide Y at sympathetic neuroeffector junctions: Existence of Y₁ and Y₂ receptors. In: Neuronal messengers in vascular function, Fernstrom Symp. No 10., pp. 231-242. Eds A. Nobin and C.H. Owman. Elsevier: Amsterdam (1987)), all of which are proposed to function as seven transmembrane-spanning G protein-coupled receptors (Dumont et al., 1992).

The Y₁ receptor recognizes NPY = PYY >> PP (Grundemar et al., 1992). The receptor requires both the N- and the C-terminal regions of the peptides for optimal recognition. Exchange of Gln³⁴ in NPY or PYY with the analogous residue from PP (Pro³⁴), however, is well-tolerated. The Y₁ receptor has been cloned from a variety of species including human, rat and mouse (Larhammar, D., A.G. Blomqvist, F. Yee, E. Jazin, H. Yoo, and C. Wahlestedt. (1992). Cloning and functional expression of a human neuropeptide Y/peptide YY receptor of the Y₁ type. J. Biol. Chem. 267: 10935-10938; Herzog, H., Y.J. Hort, H.J. Ball, G. Hayes, J. Shine, and L. Selbie. (1992). Cloned human neuropeptide Y receptor couples to two different second messenger systems. Proc. Natl. Acad. Sci. USA 89, 5794-5798; Eva, C., Oberto, A., Sprengel, R. and E. Genazzani. (1992). The murine NPY-1 receptor gene: structure and delineation of tissue specific expression. FEBS Lett. 314: 285-288; Eva, C., Keinänen, K., Monyer, H., Seeburg, P., and Sprengel, R. (1990). Molecular cloning of a novel G protein-coupled receptor that may belong to the neuropeptide receptor family. FEBS Lett. 271, 80-84). The Y₂ receptor recognizes PYY ~ NPY >> PP and is relatively tolerant of N-terminal deletion (Grundemar, L. and R. Hakanson (1994). Neuropeptide Y effector systems:

perspectives for drug development. Trends. Pharmacol. 15:153-159). The receptor has a strict requirement for structure in the C-terminus (Arg³³-Gln³⁴-Arg³⁵-Tyr³⁶-NH₂); exchange of Gln³⁴ with Pro³⁴, as in PP, is not well tolerated. The Y2 receptor has recently been cloned. The Y3 receptor is characterized by a strong preference for NPY over PYY and PP (Wahlestedt, C., Karoum, F., Jaskiw, G., Wyatt, R.J., Larhammar, D., Ekman, R., and Reis, D.J. (1991). Cocaine-induced reduction of brain neuropeptide Y synthesis dependent on medial prefrontal cortex. Proc. Natl. Acad. Sci. 88: 2978-2082). [Pro³⁴]NPY is reasonably well tolerated even though PP, which also contains Pro³⁴, does not bind well to the Y3 receptor. This receptor (Y3) has not yet been cloned. The Y4 receptor binds PP > PYY > NPY. Like the Y1, the Y4 requires both the N- and the C-terminal regions of the peptides for optimal recognition. The "atypical Y1" or "feeding" receptor is defined exclusively by injection of several pancreatic polypeptide analogs into the paraventricular nucleus of the rat hypothalamus which stimulates feeding behavior with the following rank order: NPY₂₋₃₆ ≥ NPY ~ PYY ~ [Leu³¹,Pro³⁴]NPY > NPY₁₃₋₃₆ (Kalra, S.P., Dube, M.G., Fournier, A., and Kalra, P.S. (1991). Structure-function analysis of stimulation of food intake by neuropeptide Y: Effects of receptor agonists. Physiology & Behavior 50: 5-9; Stanley, B.G., Magdalin, W., Seirafi, A., Nguyen, M.M., and Leibowitz, S.F. (1992). Evidence for neuropeptide Y mediation of eating produced by food deprivation and for a variant of the Y₁ receptor mediating this peptide's effect. Peptides 13: 581-587). The profile is similar to that of a Y1-like receptor except for the anomalous ability of NPY₂₋₃₆ to stimulate food intake with potency equivalent or better than that of NPY. A subsequent report by Balasubramaniam, A., Sheriff, S., Johnson, M.E., Prabhakaran, M., Huang, Y., Fischer, J.E., and Chance, W.T. (1994). [D-Trp³²]Neuropeptide Y: A competitive antagonist of NPY in rat hypothalamus. J. Med. Chem. 37: 311-315 showed that feeding can be regulated by [D-Trp³²]NPY. While this peptide is presented as an NPY antagonist, the published data at least in part support a stimulatory effect of [D-Trp³²]NPY on feeding. [D-Trp³²]NPY thereby represents another diagnostic tool for receptor identification.

This plasmid (pcEXV-hY5) was deposited on November 4, 1994 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and was accorded ATCC Accession No. 75943.

The plasmid which comprises the regulatory elements necessary for expression of DNA in a mammalian cell operatively linked to the DNA encoding the rat Y5 receptor as to permit expression thereof has been designated as pcEXV-rY5 (ATCC Accession No. 75944).

This plasmid (pcEXV-rY5) was deposited on November 4, 1994 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and was accorded ATCC Accession No. CRL 75944.

A method for determining whether a ligand can specifically bind to a Y5 receptor comprises contacting a cell transfected with and expressing DNA encoding the Y5 receptor with the ligand under conditions permitting binding of ligands to such receptor, detecting the presence of any such ligand specifically bound to the Y5 receptor, and thereby determining whether the ligand specifically binds to the Y5 receptor.

A method for determining whether a ligand is a Y5 receptor antagonist comprises contacting a cell transfected with and expressing DNA encoding a Y5 receptor with the ligand in the presence of a known Y5 receptor agonist, such as PYY or NPY, under conditions permitting the activation of a functional Y5 receptor response, detecting a decrease in Y5 receptor activity, and thereby determining whether the ligand is a Y5 receptor antagonist.

In an embodiment of the above-described methods, the cell is non-neuronal in origin. In a further embodiment, the non-neuronal cell is a COS-7 cell, 293 human embryonic kidney cell, NIH-3T3 cell or L-M(TK-) cell.

The cell lines are transfected with a vector which is adapted for expression in a mammalian cell which comprises the regulatory elements necessary for expression of the DNA in the mammalian cell operatively linked to the DNA encoding the mammalian Y5 receptor as to permit expression thereof.

For example, such plasmid which comprises the regulatory elements necessary for expression of DNA in a mammalian cell operatively linked to the DNA encoding the human

Y5 receptor as to permit expression thereof designated pcEXV-hY5 (ATCC Accession No. 75943).

Experimental Details

MATERIALS AND METHODS

cDNA Cloning

Total RNA was prepared by a modification of the guanidine thiocyanate method (Kingston, 1987), from 5 grams of rat hypothalamus (Rockland, Gilbertsville, PA). Poly A⁺RNA was purified with a FastTrack kit (Invitrogen Corp., San Diego, CA). Double stranded (ds) cDNA was synthesized from 7 mg of poly A⁺ RNA according to Gubler and Hoffman (Gubler, U and B.J. Hoffman. (1983). A simple and very efficient method for generating cDNA libraries. Gene. 25, 263-269), except that ligase was omitted in the second strand cDNA synthesis. The resulting DS cDNA was ligated to BstXI/EcoRI adaptors (Invitrogen Corp.), the excess of adaptors was removed by chromatography on Sephacryl 500 HR (Pharmacia®-LKB) and the ds-cDNA size selected on a Gen-Pak Fax HPLC column (Millipore Corp., Milford, MA). High molecular weight fractions were ligated in pEXJ.BS (A cDNA cloning expression vector derived from pcEXV-3; Okayama, H. and P. Berg (1983). A cDNA cloning vector that permits expression of cDNA inserts in mammalian cells. Mol. Cell. Biol. 3: 280-289; Miller, J. and Germain, R.N. (1986). Efficient cell surface expression of class II MHC molecules in the absence of associated invariant chain. J. Exp. Med. 164: 1478-1489) cut by BstXI as described by Aruffo and Seed (Aruffo, A. and Seed, B. (1987). Molecular cloning of a CD28 cDNA by a high efficiency COS cell expression system. PNAS, 84, 8573-8577). The ligated DNA was electroporated in E.Coli MC 1061 F⁺ (Gene Pulser, Biorad). A total of 3.4×10^6 independent clones with an insert mean size of 2.7 kb could be generated. The library was plated on Petri dishes (Ampicillin selection) in pools of 6.9 to 8.2×10^3 independent clones. After 18 hours amplification, the bacteria from each pool were scraped, resuspended in 4 ml of LB media and 1.5 ml processed for plasmid purification with a QIAprep-8 plasmid kit (Qiagen Inc, Chatsworth, CA). 1 ml aliquots of each bacterial pool were stored at -85°C in 20% glycerol.

Isolation of a cDNA clone encoding an atypical rat hypothalamic NPY5 receptor

DNA from pools of » 7500 independent clones was transfected into COS-7 cells by a modification of the DEAE-dextran procedure (Warden, D. and H.V. Thorne. (1968). Infectivity of polyoma virus DNA for mouse embryo cells in presence of diethylaminoethyl-dextran. *J. Gen. Virol.* 3, 371). COS-7 cells were grown in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal calf serum, 100 U/ml of penicillin, 100 mg/ml of streptomycin, 2 mM L-glutamine (DMEM-C) at 37°C in 5% CO₂. The cells were seeded one day before transfection at a density of 30,000 cells/cm² on Lab-Tek chamber slides (1 chamber, Permanox slide from Nunc Inc., Naperville, IL). On the next day, cells were washed twice with PBS, 735 µl of transfection cocktail was added containing 1/10 of the DNA from each pool and DEAE-dextran (500 mg/ml) in Opti-MEM I serum free media (Gibco®BRL LifeTechnologies Inc. Grand Island, NY). After a 30 min. incubation at 37°C, 3 µl of chloroquine (80 mM in DMEM-C) was added and the cells incubated a further 2.5 hours at 37°C. The media was aspirated from each chamber and 2 µl of 10% DMSO in DMEM-C added. After 2.5 min. incubation at room temperature, the media was aspirated, each chamber washed once with 2 ml PBS, the cells incubated 48 hours in DMEM-C and the binding assay was performed on the slides. After one wash with PBS, positive pools were identified by incubating the cells with 1 nM (3x10⁶ cpm per slide) of porcine [¹²⁵I]-PYY (NEN; SA=2200 Ci/mmol) in 20 mM Hepes-NaOH pH 7.4, CaCl₂ 1.26 mM, MgSO₄ 0.81 mM, KH₂PO₄ 0.44 mM, KCL 5.4, NaCl 10 mM, .1% BSA, 0.1% bacitracin for 1 hour at room temperature. After six washes (three seconds each) in binding buffer without ligand, the monolayers were fixed in 2.5% glutaraldehyde in PBS for five minutes, washed twice for two minutes in PBS, dehydrated in ethanol baths for two minutes each (70, 80, 95, 100%) and air dried. The slides were then dipped in 100% photoemulsion (Kodak® type NTB2) at 42°C and exposed in the dark for 48 hours at 4°C in light proof boxes containing drierite. Slides were developed for three minutes in Kodak® D19 developer (32 g/l of water), rinsed in water, fixed in Kodak® fixer for 5 minutes, rinsed in water, air dried and mounted with Aqua-Mount (Lerner Laboratories, Pittsburgh, PA). Slides were screened at 25x total magnification. A single clone, CG-18, was isolated by SIB selection as described (McCormick, 1987). DS-DNA was sequenced with a Sequenase kit (US Biochemical, Cleveland, OH) according to the manufacturer. Nucleotide and peptide sequence analysis were performed with GCG programs (Genetics Computer group, Madison, WI).

Isolation of the human Y5 homolog

Using rat oligonucleotide primers in TM 3 (sense primer; position 484-509 in SEQ ID NO:1) and in TM 6 (antisense primer; position 1219-1243 in SEQ ID NO: 1), a human hippocampal cDNA library has been screened using the polymerase chain reaction. 1 μ l (4×10^6 bacteria) of each of 450 amplified pools containing each »5000 independent clones and representing a total of 2.2×10^6 was subjected directly to 40 cycles of PCR and the resulting products analyzed by agarose gel electrophoresis. One of three positive pools was analyzed further and by sib selection a single cDNA clone was isolated and characterized. This cDNA turned out to be full length and in the correct orientation for expression. DS-DNA was sequenced with a sequenase kit (US Biochemical, Cleveland, OH) according to the manufacturer.

Cell Culture

COS-7 cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 mg/ml streptomycin) at 37°C, 5% CO₂. Stock plates of COS-7 cells were trypsinized and split 1:6 every 3-4 days. Human embryonic kidney 293 cells were grown on 150 mm plates in D-MEM with supplements (minimal essential medium) with Hanks' salts and supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 mg/ml streptomycin) at 37 °C, 5% CO₂. Stock plates of 293 cells were trypsinized and split 1:6 every 3-4 days. Mouse fibroblast LMT(k)- cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 mg/ml streptomycin) at 37 °C, 5% CO₂. Stock plates of COS-7 cells were trypsinized and split 1:10 every 3-4 days.

Stable Transfection

Human Y5 and rat Y5 receptors were co-transfected with a G-418 resistant gene into mouse fibroblast LMT(k)- cells by a calcium phosphate transfection method (Cullen, B.

(1987). Use of eukaryotic expression technology in the functional analysis of cloned genes. Methods Enzymol. 152: 685-704). Stably transfected cells were selected with G-418.

EXPERIMENTAL RESULTS

cDNA Cloning

In order to clone a rat hypothalamic "atypical" NPY receptor subtype, applicants used an expression cloning strategy in COS-7 cells (Gearing et al, 1989; Kluxen, F.W., Bruns, C. and Lubbert H. (1992). Expression cloning of a rat brain somatostatin receptor cDNA. Proc. Natl. Acad. Sci. USA 89, 4618-4622; Kieffer, B., Befort, K., Gaveriaux-Ruff, C. and Hirth, C.G. (1992). The δ -opioid receptor: Isolation of a cDNA by expression cloning and pharmacological characterization. Proc. natl. Acad. Sci. USA 89, 12048-12052). This strategy was chosen for its extreme sensitivity since it allows detection of a single "receptor positive" cell by direct microscopic autoradiography. Since the "atypical" receptor has only been described in feeding behavior studies involving injection of NPY and NPY related ligands in rat hypothalamus (see introduction), applicants first examined its binding profile by running competitive displacement studies of ^{125}I -PYY and ^{125}I -PYY₃₋₃₆ on membranes prepared from rat hypothalamus. The competitive displacement data indicate: 1) Human PP is able to displace 20% of the bound ^{125}I -PYY with an IC_{50} of 11 nM (Fig. 1 and Table 2). As can be seen in table 5, this value does not fit with the isolated rat Y1, Y2 and Y4 clones and could therefore correspond to another NPY/PYY receptor subtype. 2) [Leu₃₁, Pro₃₄] NPY (a Y1 specific ligand) is able to displace with high affinity (IC_{50} of 0.38) 27% of the bound ^{125}I -PYY₃₋₃₆ ligand (a Y2 specific ligand) (Fig. 2 and table 2). These data provide the first evidence based on a binding assay that rat hypothalamic membranes could carry an NPY receptor subtype with a mixed Y1/Y2 pharmacology (referred to as the "atypical" subtype) which fits with the pharmacology defined in feeding behavior studies.

TABLE 2: Pharmacological profile of the rat hypothalamus.

Binding data reflect competitive displacement of ^{125}I -PYY and ^{125}I -PYY₃₋₃₆ from rat hypothalamic membranes. Peptides were tested at concentrations ranging from 0.001 nM to 100 nM unless noted. The IC_{50} value corresponding to 50% displacement, and the percentage of displacement relative to that produced by 300 nM human NPY, were

determined by nonlinear regression analysis. Data shown are representative of at least two independent experiments.

TABLE 2

Peptide	IC ₅₀ Values, nM (% NPY-produced displacement)	
	¹²⁵ I-PYY	¹²⁵ I-PYY ₃₋₃₆
human NPY	0.82 (100%)	1.5 (100%)
human NPY ₂₋₃₆	2.3 (100%)	1.2 (100%)
human [Leu ³¹ ,Pro ³⁴]NPY	0.21 (44%) 340 (56%)	0.38 (27%) 250 (73%)
human PYY	1.3 (100%)	0.29 (100%)
human PP	11 (20%)	untested

Based on the above data, a rat hypothalamic cDNA library of 3×10^6 independent recombinants with a 2.7 kb average insert size was fractionated into 450 pools of »7500 independent clones. All pools were tested in a binding assay with ¹²⁵I-PYY as described (Y2 patent). Seven pools gave rise to positive cells in the screening assay (# 81, 92, 147, 246, 254, 290, 312). Since Y1, Y2, Y4 and Y5 receptor subtypes (by PCR or binding analysis) are expressed in rat hypothalamus, applicants analyzed the DNA of positive pools by PCR with rat Y1, Y2 and Y4 specific primers. Pools # 147, 246, 254 and 312 turned out to contain cDNAs encoding a Y1 receptor, pool # 290 turned out to encode a Y2 subtype, but pools # 81 and 92 were negative by PCR analysis for Y1, Y2 and Y4 and therefore likely contained a cDNA encoding a new rat hypothalamic NPY receptor (Y5). Pools # 81 and 92 later turned out to contain an identical NPY receptor cDNA. Pool 92 was subjected to sib selection as described until a single clone was isolated (designated CG-18).

The isolated clone carries a 2.8 kb cDNA. This cDNA contains an open reading frame between nucleotides 779 and 2146 that encodes a 456 amino acid protein. The long 5' untranslated region could be involved in the regulation of translation efficiency or mRNA

stability. The flanking sequence around the putative initiation codon does not conform to the Kozak consensus sequence for optimal translation initiation (Kozak, M. (1989). The scanning model for translation: an update. J. Cell Biol. 108, 229-241; Kozak, M. (1991). Structural features in eukaryotic mRNAs that modulate the initiation of translation. J. Biol. Chem. 266, 19867-19870). The hydrophobicity plot displayed seven hydrophobic, putative membrane spanning regions which makes the rat hypothalamic Y5 receptor a member of the G-protein coupled superfamily. The nucleotide and deduced amino acid sequences are shown in SEQ ID NOS: 1 and 2, respectively.

Localization studies show that the Y5 mRNA is present in several areas of the rat hippocampus. Assuming a comparable localization in human brain, applicants screened a human hippocampal cDNA library with rat oligonucleotide primers which were shown to yield a DNA band of the expected size in a PCR reaction run on human hippocampal cDNA. Using this PCR screening strategy (Gerald et al, 1994, submitted for publication), three positive pools were identified. One of these pools was analyzed further, and an isolated clone was purified by sib selection. The isolated clone (CG-19) turned out to contain a full length cDNA cloned in the correct orientation for functional expression (see below). The human Y5 nucleotide and deduced amino acid sequences are shown in SEQ ID NOS 3 and 4, respectively. When compared to the rat Y5 receptor the human sequence shows 84.1% nucleotide identity and 87.2% amino acid identity. The rat protein sequence is one amino acid longer at the very end of both amino and carboxy tails of the receptor when compared to the rat. Both pharmacological profiles and functional characteristics of the rat and human Y5 receptor subtype homologs may be expected to match closely.

When the human and rat Y5 receptor sequences were compared to other NPY receptor subtypes or to other human G protein-coupled receptor subtypes, both overall and transmembrane domain identities are very low, showing that the Y5 receptor genes are not closely related to any other previously characterized cDNAs.

The compounds according to the present invention and their pharmaceutically acceptable salts have proven to exhibit pronounced and selective affinity to the Y5 receptor subtype (shown in Y5 binding test) and in vitro and in vivo antagonistic properties. These properties are shown in vitro by their ability to inhibit NPY-induced calcium increase in stable transfected cells expressing the Y5 receptor and in vivo by their ability to inhibit food intake

induced by intracerebroventricular application of NPY or 24 h food deprivation in conscious rats.

Binding experiments

The selective affinity of the compounds according to the present invention to the Y5 receptor is detected in a Y5 binding assay using LM(tk-)-h-NPY5-7 cells which stably express the human NPY Y5 receptor or HEK-293 cells stably expressing the rat NPY Y5 receptor.

The following buffers are used for the preparation of membranes and for binding assay:

a) buffer 1 (homogenisation buffer, pH 7.7 at 4°C) contains Tris-HCl [FLUKA, Buchs, Switzerland] (20 mM) and ethylenediamine tetraacetate (EDTA) [FLUKA, Buchs, Switzerland] (5 mM); b) buffer 2 (suspension buffer, pH: 7.4 at room temperature) contains N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) [Boehringer Mannheim, Germany] (20 mM), NaCl (10 mM), CaCl₂ (1.26 mM), MgSO₄ (0.81 mM) and KH₂PO₄ (0.22 mM); buffer 3 (binding buffer, pH 7.4 at room temperature) contains HEPES (20 mM), NaCl (10 mM), CaCl₂ (1.26 mM), MgSO₄ (0.81 mM), KH₂PO₄ (0.22 mM) and 1 mg/ml bovine serum albumin [FLUKA].

Cells are washed in phosphate buffered saline and harvested using a rubber policeman. The cells are homogenised using a Polytron homogeniser (3 bursts of 8 seconds) in ice-cold hypotonic buffer (buffer 1, pH 7.7 at 4°C). The homogenate is centrifuged at 32,000 x g for 20 min at 4°C. The pellets are resuspended in the same buffer and recentrifuged. The final pellets are suspended in buffer 2. Protein concentration is measured by the method of Bradford using the Pierce reagent [PIERCE, Rockford, USA], with bovine serum albumin as standard. The crude membrane preparation is aliquoted, flash-frozen in liquid nitrogen and stored at -80°C. Before use, 0.1% (1 mg/ml) bovine serum albumin is added.

¹²⁵I-[Pro³⁴]hPYY (60 pM, Anawa, Wangen, Switzerland) dissolved in buffer 3 is used as radioligand. All test compounds are dissolved in dimethyl sulfoxide (DMSO) at 10⁻² M and diluted to 10⁻³ M in buffer 3. Subsequent dilutions are in buffer 3 plus 10% DMSO. Incubations are performed in Millipore Multiscreen FC filter plates [Millipore, Bedford, USA]. The filters in each well are pretreated with 2% polyethyleneimine for 30 min and rinsed once with 300 microL buffer 3 before use. The following are pipetted into each well: 20 microL buffer 3, 25 microL ¹²⁵I-[Pro³⁴]hPYY [SAXON, Hannover, Germany] (600 pM); 25 microL test compound (or binding buffer for the controls); 180 microL crude membrane suspension (approximately 5 microg protein). Incubations are performed at room temperature for 2h.

Non-specific binding is defined as the binding remaining in the presence of 1 microM [Pro³⁴]hPYY. The incubations are terminated by rapid filtration and washing four times with 300microL phosphate buffered saline. The filters are removed from the wells, placed into plastic tubes and assayed for radioactivity in a gamma counter [Gammamaster, WALLAC, Finland].

The IC₅₀ values of the compounds according to this invention at the human Y5 receptor range especially between about 0.1 nM and about 10 microM.

Measurements of calcium transient

For the determination of in vitro antagonistic properties of the compounds according to the present invention, stably transfected LM(tk-)-hY5-7 cells are used in which a NPY-induced calcium transient is measured as described below. Cells are harvested in a medium containing EDTA (0.5 mM) and phosphate buffered saline (PBS). Cells are then washed in phosphate buffered saline solution and loaded for 90 min at room temperature and pH 7.4 with 10 microM FLUO-AM (fluoro-3-acetoxy methylester, supplemented with pluronic acid as suggested by the manufacturer, Molecular Probes Inc., Eugene, Oregon, USA) in a cell culture buffer of the following composition (NaCl 120 mM, MgCl₂ 1 mM, KCl 5.4 mM, NaH₂PO₄ 0.33 mM, glucose 11 mM, taurine 5 mM, pyruvate 2 mM, glutamine 1.5 mM HEPES 10 mM, insulin 10 U/l, BSA 0.1% at for 90 min at room temperature. After centrifugation the cells are resuspended in the cell culture buffer at a concentration of 3-4 million cells/ml and supplemented with 200 microM sulfinpyrazone.

Calcium transients are measured at room temperature in a milliliter plate using a Cytofluor 2350 (Millipore) with wavelength settings at 485 nm for excitation and 530 nm for emission. 180 microL of cells suspension are preincubated in the presence of various amounts of compounds dissolved in 2 microL DMSO in triplicates (or 2 microL DMSO for the controls) for 5 min and then NPY is added at a final concentration of 100 nM. The compound concentrations giving 50% inhibition of the maximum of the Ca transients are then calculated.

In this cell system, NPY induces Ca transients with an EC₅₀ of 50 nM. The data are analyzed using a Microsoft Excel software. The concentrations which cause a 50% inhibition of the initial control values are given as IC₅₀ values. The IC₅₀ values are determined for the compounds according to the present invention and their pharmaceutically acceptable salts.

The property of the compounds according to the present invention and their pharmaceutically acceptable salts to inhibit NPY-induced increase intracellular calcium indicates their antagonistic properties with IC₅₀ values ranging especially between about 0.1 nM and about 10 microM. Representatives are, for example, the final products of working examples 1 and 2, for which following IC₅₀ values [μ M/L] were determined: 0.008 (Ex. 1); 0.01 (Ex. 2).

Measurements of NPY-induced food intake in conscious rats

In addition this antagonistic property of the Y₅ receptor subtype is also observed in-vivo in conscious rats by their ability to inhibit NPY-induced food intake. For these determinations food intake is measured in normal satiated rats after intracerebroventricular application (i.c.v.) of neuropeptide Y [BACHEM, Feinchemikalien, Bubendorf, Switzerland] in the presence or absence of the compounds according to the present invention. Male Sprague-Dawley rats weighing 180-220 g are used for all experiments. They are individually housed in stainless steel cages and maintained on a 11:13 h light-dark schedule (lights off at 1800 h) under controlled temperature (21-23 °C) at all times. Water and food (NAFAG lab chow pellets) [NAFAG, Gossau, Switzerland] are available ad libitum.

Under pentobarbital [VETERINARIA AB, Zürich, Switzerland] anesthesia, all rats are implanted with a stainless steel guide cannula targeted at the right lateral ventricle. Stereotaxic coordinates, with the incisor bar set -2.0 mm below interaural line, are : -0.8 mm anterior and +1.3 mm lateral to bregma. The guide cannula is placed on the dura. Injection cannulas extended the guide cannulas -3.8 mm ventrally to the skull surface. Animals are allowed at least 4 days of recovery postoperatively before being used in the experiments. Cannula placement is checked postoperatively by testing all rats for their drinking response to a 50 ng intracerebroventricular (icv) injection of angiotensin II . Only rats which drink at least 2.5 ml of water within 30 min after angiotensin II injection are used in the feeding studies. Injections are made in the morning 2 hours after light onset. Peptides are injected in artificial cerebrospinal fluid (ACSF) [FLUKA, Buchs, Switzerland] in a volume of 5 μ l. The ACSF contains NaCl 124 mM, KCl 3.75 mM, CaCl₂ 2.5 mM, MgSO₄ 2.0 mM, KH₄PO₄ 0.22 mM, NaHCO₃ 26 mM and glucose 10 mM. NPY (300 pmole) is administered by the intracerebroventricular route 10-60 minutes after administration of compounds or vehicle DMSO/water (10%,v/v) or cremophor/water (20%,v/v) [SIGMA, Buchs, Switzerland].

Food intake is measured by placing preweighed pellets into the cages at the time of NPY injection. Pellets are removed from the cage subsequently at each time point indicated in the figures and replaced with a new set of preweighed pellets.

All results are presented as means \pm SEM. Statistical analysis is performed by analysis of variance using Student-Newman-Keuls test.

The compounds according to the present invention inhibit NPY-induced food intake in rats in a range especially of about 0.01 to about 100 mg/kg after oral, intraperitoneal, subcutaneous or intravenous administration.

Measurements of food intake in 24 hours food deprived rats

Based on the observation that food deprivation induces an increase in the hypothalamic NPY levels, it is assumed that NPY mediates food intake induced by food deprivation. Thus, the compounds according to the present invention are also tested in rats after 24 hours food deprivation. These experiments are conducted with male Sprague-Dawley (CIBA-GEIGY AG, Sisseln, Switzerland) rats weighing between 220 and 250 g. The animals are housed in individual cages for the duration of the study and allowed free access to normal food together with tap water. The animals are maintained in room with a 12 h light/dark cycle (8 a.m. to 8.00 p.m. light) at 24°C and monitored humidity. After placement into the individual cages the rats undergo a 2-4 days equilibration period, during which they are habituated to their new environment and to eating a powdered or pellet diet [NAFAG, Gossau, Switzerland]. At the end of the equilibration period, food is removed from the animals for 24 hours starting at 8.00 a.m. At the end of the fasting period the animals are injected intraperitoneally, intravenously or orally either with the compounds according to the present invention or an equivalent volume of vehicle DMSO/water (10%, v/v) or cremophor/water (20%, v/v) and 10-60 min later the food is returned to them. Food intake at various time periods is monitored over the following 24 hour period. Inhibition of food intake by the compounds according to the present invention is given in percentage of the respective control vehicle-treated rats.

The compounds according to the present invention inhibit food intake in this food deprived rat model in a range especially of about 0.01 to about 100 mg/kg after oral, intraperitoneal, subcutaneous or intravenous administration. Representatives are, for example, the final products of working examples 3 and 4, for which an inhibition of food intake of 58% or 55%,

respectively, versus the respective control vehicle-treated animals after i.p. application of 30 mg/kg was determined.

Measurements of food intake in obese Zucker rats

The antiobesity efficacy of the compounds according to the present invention can also be shown in Zucker obese rats, an art-known animal model of obesity. These studies are conducted with male Zucker fatty rats (fa/fa) [HARLAN CPB, Austerlitz, NL] weighing between 480 and 500 g. Animals are individually housed in metabolism cages for the duration of the study and allowed free access to powdered food together with tap water. The animals are maintained in a room with a 12 hour light/dark cycle (8 a.m. to 8.00 p.m. light) at 24°C and monitored humidity. After placement into the metabolism cages the rats undergo a 6 day equilibration period, during which they are habituated to their new environment and to eating a powdered diet. At the end of the equilibration period, food intake during the light and dark phases is determined. After a 3 day control period, the animals are treated with the compounds according to the present invention or vehicle DMSO/water (10%, v/v) or cremophor/water (20%, v/v).

The compounds according to the present invention inhibit food intake in Zucker obese rats in a range especially of about 0.01 to about 100 mg/kg after oral, intraperitoneal, subcutaneous or intravenous administration.

The above experiments clearly demonstrate that the Y5 receptor subtype is the primary mediator of NPY-induced feeding and that corresponding antagonists can be used for the treatment of obesity and related disorders [*Nature*, Vol. 382, 168-171 (1996)].

The compounds according to the present invention can inhibit food intake induced either by intracerebroventricular application of NPY or by food deprivation or as well as spontaneous eating in the Zucker obese rat. Thus, the compounds according to the present invention can especially be used for the prophylaxis and treatment of disorders or diseases associated with the Y5 receptor subtype, especially in the treatment of disorders or disease states in which the NPY-Y5 receptor subtype is involved, preferably, in the treatment of diseases caused by eating disorders, such as obesity, bulimia nervosa, diabetes, dyslipidemia, and hypertension, furthermore in the treatment of memory loss, epileptic seizures, migraine, sleep disturbance, and pain and additionally in the treatment of sexual/reproductive

disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion and diarrhea.

The compounds according to the present invention act as antagonists of neuropeptide Y (NPY) binding at the Y5 receptor subtype. By virtue of their Y5 receptor antagonistic property, the compounds of the formula (I) and their pharmaceutically acceptable salts can therefore be used, for example, as pharmaceutical active ingredients in pharmaceutical compositions which are employed, for example, for the prophylaxis and treatment of diseases and disorders associated with NPY Y5 receptor subtype, especially in the treatment of disorders or disease states in which the NPY-Y5 receptor subtype is involved, preferably, in the treatment of diseases caused by eating disorders, such as obesity, bulimia nervosa, diabetes, dyslipidemia, and hypertension, furthermore in the treatment of memory loss, epileptic seizures, migraine, sleep disturbance, and pain, and additionally in the treatment of sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion and diarrhea.

The invention relates to the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as described hereinbefore and hereinafter for the manufacture of a pharmaceutical composition for the prophylaxis and treatment of diseases or disorders associated with NPY Y5 receptor subtype, especially in the treatment of disorders or disease states in which the NPY-Y5 receptor subtype is involved, preferably, in the treatment of diseases caused by eating disorders, such as obesity, bulimia nervosa, diabetes, dyslipidemia, and hypertension, furthermore in the treatment of memory loss, epileptic seizures, migraine, sleep disturbance, and pain, and additionally in the treatment of sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion and diarrhea.

The invention relates to a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as described hereinbefore and hereinafter for the treatment of diseases or disorders associated with NPY Y5 receptor subtype, preferably, in the prophylaxis and treatment of diseases caused by eating disorders, such as obesity, bulimia nervosa, diabetes, dyslipidemia, and hypertension, furthermore in the treatment of memory loss, epileptic seizures, migraine, sleep disturbance, and pain, and

additionally in the treatment of sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion and diarrhea.

The invention relates especially to a method of treatment and prophylaxis of disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

- (i) hydrogen, halogen, nitro, cyano, lower alkyl, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by lower alkoxy, by substituted amino, by lower alkoxy-carbonyl, or by N-substituted carbamoyl;
- (ii) substituted amino;
- (iii) hydroxy, lower alkoxy, lower alkoxy-lower alkoxy, C₃-C₈-cycloalkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, lower alkoxy-carbonyl-oxy, or N-substituted aminocarbonyl-oxy;
- (iv) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (v) N-substituted carbamoyl;
- (vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-O-R, -NR₀₁-CO-R, -NR₀₁-CO-NR₁-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R being as defined below and R₀₁ being as defined above, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O, S(O)_n or NR₀} or which is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring]; or
- (vii) an element of formula -X₁(X₂)(X₃) wherein, (a) if X₁ is -CH-, X₂ together with X₃ represent a structural element of formula -X₄-(CO)_p-(CH₂)_o-, -(CH₂)_q-X₄-(CO)_p-(CH₂)_r-, or -(CH₂)_s-X₄-CO-(CH₂)_t-; or, (b) if X₁ is -N-, X₂ together with X₃ represent a structural element of formula -CO-(CH₂)_u-; [X₄ being -CH₂-, -N(R₀₁)- or -O-; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X₄ is different from -CH₂-];

R₂ and R₃, independently of one another, represent

- (i) hydrogen, lower alkyl, lower alkenyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, N-substituted carbamoyl, and -S(O)_n-R;

R₂ and R₃ together represent lower alkylene [which may be interrupted by O, S(O)_n, or NR₀] or represent lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring;

X represents N or CH; and the integer n is 0 or 1;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

- (i) halogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, nitro, cyano;
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryloxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iv) amino, substituted amino;
- (v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, is derived and selected from the group consisting of phenyl, biphenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furyl, thienyl,

pyridyl, indolyl, indazolyl, benzofuryl, benzothiophenyl, benzimidazolyl, quinoliny, isoquinoliny, or quinazoliny;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, S(O)_n or NR₀] or is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring, or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen or lower alkyl;

wherein, in each case, R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy.

The invention relates especially to a method of treatment and prophylaxis of disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

- (i) hydrogen, halogen, nitro, cyano, lower alkyl, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryl, or lower alkyl which is substituted by halogen, by lower alkoxy, by substituted amino, by lower alkoxy-carbonyl, or by substituted carbamoyl;
- (ii) substituted amino;
- (iii) hydroxy, lower alkoxy, C₃-C₈-cycloalkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, or lower alkoxy-lower alkoxy;
- (iv) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (v) N-substituted carbamoyl;
- (vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R and R₁ being as defined below, or the

group $-N(R)(R_{01})$ represents amino which is di-substituted by lower alkylene {which may be interrupted by O or NR_0 } or which is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring]; or

(vii) an element of formula $-X_1(X_2)(X_3)$ wherein, (a) if X_1 is $-CH-$, X_2 together with X_3 represent a structural element of formula $-X_4-(CO)_p-(CH_2)_o-$, $-(CH_2)_q-X_4-(CO)_p-(CH_2)_r-$, or $-(CH_2)_s-X_4-CO-(CH_2)_t-$; or, (b) if X_1 is $-N-$, X_2 together with X_3 represent a structural element of formula $-CO-(CH_2)_u-$; [X_4 being $-CH_2-$, $-N(R_{01})-$ or $-O-$; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X_4 is different from $-CH_2-$];

R_2 and R_3 , independently of one another, represent

(i) hydrogen, lower alkyl, lower alkenyl, (carbocyclic or heterocyclic) aryl, or (carbocyclic or heterocyclic) aryl-lower alkyl;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: lower alkoxy, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, and substituted carbamoyl;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O, $S(O)_n$, or NR_0];

X represents N or CH; and the integer n is 0 or 1;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl-lower alkyl, lower alkoxy, lower alkenyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C_3 - C_8 -cycloalkyl, (carbocyclic or heterocyclic) aryloxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

- (iv) substituted amino;
- (v) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, is derived from phenyl, naphthyl or pyridyl;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, S(O)_n or NR₀] or is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring, or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen or lower alkyl;

wherein, in each case, R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy.

The invention relates especially to a method of treatment and prophylaxis of disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

- (i) hydrogen, halogen, cyano, nitro, lower alkyl, C₃-C₈-cycloalkyl, or phenyl;
- (ii) amino which is mono-substituted by lower alkyl, phenyl or pyridyl, or which is disubstituted by lower alkyl or by C₂-C₆-alkylene;
- (iii) hydroxy or lower alkoxy which is unsubstituted or substituted by C₃-C₈-cycloalkyl or by phenyl;
- (iv) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, or -SO₂-NR₀₁-R, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or

naphthyl, R_{01} being hydrogen or lower alkyl, or the group $-N(R)(R_{01})$ represents amino which is mono-substituted by lower alkyl, by hydroxy-lower alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C_2-C_6 -alkylene {which may be interrupted by O or NR_0 , R_0 being hydrogen or lower alkyl}};

R_2 represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

R_3 represents hydrogen;

X represents N or CH; and the integer n is 0 or 1;

wherein any aryl moiety, if not designated otherwise, and the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, lower alkyl, halo-alkyl, lower alkoxy, hydroxy, hydroxy-lower alkoxy, and lower alkoxy-lower alkoxy.

The invention relates especially to a method of treatment of and prophylaxis disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R_1 represents (i) hydrogen, halogen, cyano, nitro, lower alkyl, C_3-C_8 -cycloalkyl, or phenyl;

(ii) amino which is mono-substituted by lower alkyl, phenyl or pyridyl, or which is disubstituted by lower alkyl or by C_2-C_6 -alkylene;

(iii) hydroxy or lower alkoxy which is unsubstituted or substituted by C_3-C_8 -cycloalkyl, or by phenyl;

(iv) a group selected from $-NR_{01}-CO-R$, $-NR_{01}-SO_2-R$, $-NR_{01}-SO_2-NR_{01}-R$, $-SO_2-R$, or $-SO_2-NR_{01}-R$, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or naphthyl, R_{01} being hydrogen or lower alkyl, or the group $-N(R)(R_{01})$ represents amino which is mono-substituted by lower alkyl, by hydroxy-lower alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C_2-C_6 -alkylene {which may be interrupted by O or NR_0 , R_0 being hydrogen or lower alkyl}};

R_2 represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

- 39 -

R₃ represents hydrogen;

X represents N or CH; and the integer n is 0 or 1;

wherein any aryl moiety, if not designated otherwise, and the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, lower alkoxy, hydroxy, and hydroxy-lower alkoxy.

The invention relates especially to a method of treatment of and prophylaxis disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents hydrogen, amino which is disubstituted by by C₂-C₆-alkylene, especially pentylene, or C₁-C₄-alkoxy, especially methoxy; or a group selected from -NH-SO₂-R, -SO₂-R, or -SO₂-NH-R, [R being C₁-C₄-alkyl, or naphthyl, or the group -NH(R) represents amino which is mono-substituted by C₁-C₄-alkyl, by hydroxy-C₁-C₄-alkyl, or by naphthyl, or which is di-substituted by C₁-C₄-alkyl or by C₂-C₆-alkylene {which may be interrupted by O or NR₀, R₀ being hydrogen or C₁-C₄-alkyl}];

and, in each case,

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, and oxy-C₁-C₄-alkylene-oxy; and

R₃ represents hydrogen;

X represents N or CH; and the integer n is 0 or 1;

wherein the benzo ring A is unsubstituted or substituted by C₁-C₄-alkoxy.

The invention relates especially to a method of treatment of and prophylaxis disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents -(CH₂)_n- and the integer n is 1-3;

R₁ represents the group of formula NH-SO₂-R and R is naphthyl;

R₂ and R₃ each are hydrogen;

X is N or CH; and

the ring A is unsubstituted or substituted by C₁-C₄-alkoxy, especially methoxy.

The invention relates especially to a method of treatment of and prophylaxis disorders and diseases associated with NPY receptor subtype Y5 comprising administering to a warm-blooded animal in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which alk represents $-(CH_2)_o-$ and the integer o is 1 or 2; R₁ represents the group of formula NH-SO₂-R and R is naphthyl; R₂ and R₃ each are hydrogen; X is CH; the integer n is 1; and the ring A is unsubstituted.

The invention likewise relates to a new compound of formula (I) or a salt thereof as described hereinbefore or hereinafter.

The invention relates especially to a new compound of formula (I) or a salt thereof, e.g. in which

alk represents a single bond or lower alkylene;

the integer n is 0 or 1;

R₁ represents

(vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-O-R, -NR₀₁-CO-R, -NR₀₁-CO-NR₀₁-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R and R₁ being as defined below, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O, S(O)_n or NR₀ or which may be condensed at two adjacent carbon atoms with a benzene ring}]; or

(vii) an element of formula -X₁(X₂)(X₃) wherein, (a) if X₁ is -CH-, X₂ together with X₃ represent a structural element of formula -X₄-(CO)_p-(CH₂)_o-, -(CH₂)_q-X₄-(CO)_p-(CH₂)_r-, or -(CH₂)_s-X₄-CO-(CH₂)_t-; or, (b) if X₁ is -N-, X₂ together with X₃ represent a structural element of formula -CO-(CH₂)_u- [X₄ being -CH₂-, -N(R₀₁)- or -O-; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X₄ is different from -CH₂-];

R₂ and R₃, independently of one another, represent

- (i) hydrogen, lower alkyl, lower alkenyl, lower alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino,

substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, N-substituted carbamoyl, and $-S(O)_n-R$;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O, $S(O)_n$, NR_0 or which may be condensed at two adjacent carbon atoms with a benzene ring];

X represents N or CH;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, lower alkenyl, lower alkynyl, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, lower alkynyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, (carbocyclic or heterocyclic) aroyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iv) amino, substituted amino;

(v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, the substituted amino group of substituted amino, N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C_3-C_8 -cycloalkyl, by C_3-C_8 -cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, $S(O)_n$ or NR_0 or which may be condensed at two adjacent carbon atoms with a benzene

ring] , or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by $-\text{CO}-(\text{O})_v-\text{R}$ and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R_0 represents hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower alkanoyl, (carbocyclic or heterocyclic) aroyl, $-\text{SO}_2-\text{R}$, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;

wherein, in each case, R_{01} represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy; or a salt thereof; and the use of compounds of formula (I) or a salt thereof,

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents a single bond or lower alkylene;

R_1 represents

(vi) a group selected from $-\text{CH}(\text{OH})-\text{R}$, $-\text{CO}-\text{R}$, $-\text{NR}_{01}-\text{CO}-\text{O}-\text{R}$, $-\text{NR}_{01}-\text{CO}-\text{R}$, $-\text{NR}_1-\text{CO}-\text{NR}_{01}-\text{R}$, $-\text{NR}_{01}-\text{SO}_2-\text{R}$, $-\text{NR}_{01}-\text{SO}_2-\text{NR}_{01}-\text{R}$, $-\text{SO}_2-\text{R}$, $-\text{SO}_2-\text{NR}_{01}-\text{R}$, or $-\text{SO}_2-\text{NR}_{01}-\text{CO}-\text{R}$, [R and R_1 being as defined below, or the group $-\text{N}(\text{R})(\text{R}_{01})$ represents amino which is di-substituted by lower alkylene {which may be interrupted by O , $\text{S}(\text{O})_n$ or NR_0 } or which is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring]; or
 (vii) an element of formula $-\text{X}_1(\text{X}_2)(\text{X}_3)$ wherein, (a) if X_1 is $-\text{CH}-$, X_2 together with X_3 represent a structural element of formula $-\text{X}_4(\text{CO})_p(\text{CH}_2)_o-$, $-(\text{CH}_2)_q-\text{X}_4(\text{CO})_p(\text{CH}_2)_r-$, or $-(\text{CH}_2)_s-\text{X}_4-\text{CO}(\text{CH}_2)_t-$; or, (b) if X_1 is $-\text{N}-$, X_2 together with X_3 represent a structural element of formula $-\text{CO}(\text{CH}_2)_u-$; [X_4 being $-\text{CH}_2-$, $-\text{N}(\text{R}_{01})-$ or $-\text{O}-$; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X_4 is different from $-\text{CH}_2-$];

R_2 and R_3 , independently of one another, represent

(i) hydrogen, lower alkyl, lower alkenyl, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
 (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, substituted carbamoyl, and $-\text{S}(\text{O})_n-\text{R}$;

R₂ and R₃ together represent lower alkylene [which may be interrupted by O, S(O)_n, or NR₀] or represent lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring;

X represents N or CH; and the integer n is 0 or 1;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryloxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iv) amino, substituted amino;

(v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, is derived and selected from the group consisting of phenyl, biphenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furyl, thienyl, pyridyl, indolyl, indazolyl, benzofuryl, benzothiophenyl, benzimidazolyl, quinolinyl, isoquinolinyl, or quinazolinyl;

wherein, in each case, the amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O,

$S(O)_n$ or NR_0] or is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring, or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by $-CO-(O)_v-R$ and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R_0 represents hydrogen or lower alkyl;

wherein, in each case, R_{01} represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents a single bond or C_1 - C_3 -alkylene;

R_1 represents

(iv) a group selected from $-NR_{01}-CO-R$, $-NR_{01}-SO_2-R$, $-NR_{01}-SO_2-NR_{01}-R$, $-SO_2-R$, or $-SO_2-NR_{01}-R$, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or naphthyl, R_{01} being hydrogen or lower alkyl, or the group $-N(R)(R_{01})$ represents amino which is mono-substituted by lower alkyl, by hydroxy-lower alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C_2 - C_6 -alkylene {which may be interrupted by O or NR_0 , R_0 being hydrogen or lower alkyl}];

R_2 represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

R_3 represents hydrogen;

X represents N or CH ;

wherein the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, lower alkoxy, hydroxy, hydroxy-lower alkoxy, and lower alkoxy-lower alkoxy.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents a single bond or C_1 - C_3 -alkylene; R_1 represents a group selected from $-NR_{01}-CO-R$, $-NR_{01}-SO_2-R$, $-NR_{01}-SO_2-NR_{01}-R$, $-SO_2-R$, or $-SO_2-NR_{01}-R$, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or naphthyl, R_{01} being hydrogen or lower alkyl; or the group $-N(R)(R_{01})$ represents amino which is mono-substituted by lower alkyl, by hydroxy-lower

alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C₂-C₆-alkylene {which may be interrupted by O or NR₀, R₀ being hydrogen or lower alkyl}};

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

R₃ represents hydrogen;

X represents N or CH;

wherein the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, lower alkoxy, hydroxy, hydroxy-lower alkoxy, and lower alkoxy-lower alkoxy.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents a single bond or C₁-C₃-alkylene;

R₁ represents (iv) a group selected from -NH-SO₂-R, -SO₂-R, or -SO₂-NH-R, [R being C₁-C₄-alkyl, or naphthyl, or the group -NH(R) represents amino which is mono-substituted by C₁-C₄-alkyl, by hydroxy-C₁-C₄-alkyl, or by naphthyl, or which is di-substituted by C₁-C₄-alkyl or by C₂-C₆-alkylene {which may be interrupted by O or NR₀, R₀ being hydrogen or C₁-C₄-alkyl}};

and, in each case,

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, and oxy-C₁-C₄-alkylene-oxy; and

R₃ represents hydrogen;

X represents N or CH;

wherein the benzo ring A is unsubstituted or substituted by C₁-C₄-alkoxy.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents a single bond or C₁- or C₂-alkylene; and R₁ represents -SO₂-R or -SO₂-NH-R and R being naphthyl, especially 1- or 2-naphthyl; and, in each case,

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by lower alkoxy;

R₃ represents hydrogen; and

X represents CH;

wherein the benzo ring A is unsubstituted or substituted by C₁-C₄-alkoxy, especially, methoxy, preferably in position 8 of the quinazoline ring.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents methylene; and R₁ represents -SO₂-NH-R and R being naphthyl, especially 1- or 2-naphthyl; and, in each case,

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by lower alkoxy;

R₃ represents hydrogen; and

X represents CH;

wherein the benzo ring A is unsubstituted or substituted by C₁-C₄-alkoxy, especially, methoxy, preferably in position 8 of the quinazoline ring.

The invention relates especially to a new compound of formula (I) or a salt thereof in which alk represents methylene; and R₁ represents -SO₂-NH-R and R being naphthyl, especially 1- or 2-naphthyl; and, in each case,

R₂ represents hydrogen;

R₃ represents hydrogen; and


X represents CH;

wherein the benzo ring A is unsubstituted or substituted by C₁-C₄-alkoxy, especially, methoxy, preferably in position 8 of the quinazoline ring.

The invention relates in particular to the novel compounds shown in the examples and to the modes of preparation described therein.

The invention relates to processes for the preparation of the compounds according to the invention. The preparation of new compounds of the formula (I) and their salts comprises, for example,
(a) reacting a compound of formula (IIa) or a salt thereof

- 47 -




in which Z_1 represents a leaving group,
with a compound of formula (IIb) or a salt or tautomer thereof



or a salt thereof; or

(b) reacting a compound of formula (IIIa) or a salt thereof



in which Z_2 is a leaving group,
with a compound of formula $\text{HN}(\text{R}_2)(\text{R}_3)$ (IIIb) or a salt thereof,
and, if desired, converting a compound (I) obtainable according to the process or
in another manner, in free form or in salt form, into another compound (I),
separating a mixture of isomers obtainable according to the process and isolating
the desired isomer and/or converting a free compound (I) obtainable according to
the process into a salt or converting a salt of a compound (I) obtainable according
to the process into the free compound (I) or into another salt.

The reactions described above and below in the variants are carried out in a
manner known per se, for example in the absence or, customarily, in the

presence of a suitable solvent or diluent or a mixture thereof, the reaction, as required, being carried out with cooling, at room temperature or with warming, for example in a temperature range from about -80°C up to the boiling point of the reaction medium, preferably from about -10° to about $+200^{\circ}\text{C}$, and, if necessary, in a closed vessel, under pressure, in an inert gas atmosphere and/or under anhydrous conditions. The person skilled in the pertinent art is especially referred to the methods as outlined in the working examples based upon which the person skilled in the art is enabled to carry out the manufacture of the compounds of formula (I).

Salts of starting materials which have at least one basic centre, for example of the formula IIIb, are appropriate acid addition salts, while salts of starting materials which have an acidic group, for example of the formula (IIb), are present as salts with bases, in each case as mentioned above in connection with corresponding salts of the formula (I).

A leaving group Z_1 or Z_2 , respectively, is, for example, reactive esterified hydroxy, or is $\text{R}'\text{-S(O)}_p\text{-}$ [the integer u being 0, 1 or 2 and R' being lower alkyl, halo-lower alkyl or aryl, such as methyl, trifluoromethyl or p-toluy], or is lower alkoxy.

Reactive esterified hydroxyl Z_4 is in particular hydroxyl esterified with a strong inorganic acid or organic sulfonic acid, for example halogen, such as chlorine, bromine or iodine, sulfonyloxy, such as hydroxysulfonyloxy, halosulfonyloxy, for example fluorosulfonyloxy, $\text{C}_1\text{-C}_7\text{-alkane-sulfonyloxy}$ which is unsubstituted or substituted, for example by halogen, for example methane- or trifluoromethanesulfonyloxy, $\text{C}_5\text{-C}_7\text{-cycloalkanesulfonyloxy}$, for example cyclohexanesulfonyloxy, or benzenesulfonyloxy which is unsubstituted or substituted, for example by $\text{C}_1\text{-C}_7\text{-alkyl}$ or halogen, for example p-bromobenzene- or p-toluenesulfonyloxy. Preferred Z_1 or Z_2 is chloro, bromo or iodo, methanesulfonyloxy or trifluoromethanesulfonyloxy, or p-toluenesulfonyloxy, or methylthio or methoxy.

The reactions of process variants (a) and (b) are carried out, if necessary, in the presence of a base. Suitable bases are, for example, alkali metal hydroxides, hydrides, amides, alkanolates, carbonates, triphenylmethylenes, di-lower


alkylamides, aminoalkylamides or lower alkylsilylamides, naphthaleneamines, lower alkylamines, basic heterocycles, ammonium hydroxides, and carbocyclic amines. Examples which may be mentioned are sodium hydroxide, sodium hydride, sodium amide, sodium methoxide, sodium ethoxide, potassium tert-butoxide, potassium carbonate, lithium triphenylmethyllide, lithium diisopropylamide, potassium 3-(aminopropyl)amide, potassium bis(trimethylsilyl)amide, dimethylaminonaphthalene, di- or triethylamine, or ethyldiisopropylamine, N-methylpiperidine, pyridine, benzyltrimethylammonium hydroxide, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).

The starting material of formulae (IIa), (IIb), (IIIa), and (IIIb) is essentially known or is accessible analogously to preparation processes known per se.

The starting material of formula (IIb) in which R_2 represents N-acylated or N-alkylated amino, such as a group of formula $-NR_1-CO-O-R$, $-NR_1-CO-R$, $-NR_1-CO-NR_1-R$, $-NR_1-SO_2-R$, $-NR_1-SO_2-NR_1-R$, or N-substituted amino, is accessible, for example, by N-acylating or by N-alkylating, respectively, a, preferably N-protected, compound of the formula $NH(R_1)-alk_1-X-alk_2-Z_3$ (IIc) in which Z_3 represents a group which is convertible to R_2 , such as amino, carboxy, or hydroxy. Conventional protecting groups may be used, for example, t-butoxycarbonyl which will be split off after the N-acylation or the N-alkylation, respectively. The starting material of formula (IIb) in which R_2 represents carbamoyl or N-substituted carbamoyl, or esterified carboxy, can be manufactured starting from a compound of formula (IIc) in which Z_3 represents carboxy. The esterification or amidation can be carried out in a manner known per se. Starting from a compound of formula (IIc) in which Z_3 is hydroxy, corresponding etherified or esterified derivatives are accessible using etherification or esterification methods known in the art.

The starting material of formula (IIIa) is accessible, for example, by selectively converting the 4- Z_2 -group into a group which is deactivated, for example, by selectively hydrolyzing a compound of formula (IIIc)

- 50 -



or a salt thereof to form a corresponding 4-hydroxy-compound which is in the next step reacted with a compound of formula (IIb) to introduce the corresponding side chain into position 2 of the quinazolin ring. Reactivation of the 4-position, for example, by reaction with a halogenating agent, such as POCl_3 , leads to corresponding compounds of formula (IIIa).

A compound according to the invention which is obtainable by the process can be converted into another compound according to the invention in a manner known per se.

A compound according to the invention containing hydroxyl can be etherified by methods known per se. The etherification can be carried out, for example, using an alcohol, such as a substituted or unsubstituted lower alkanol, or a reactive ester thereof. Suitable reactive esters of the desired alcohols are, for example, those with strong inorganic or organic acids, such as corresponding halides, sulfates, lower alkanesulfonates or substituted or unsubstituted benzenesulfonates, for example chlorides, bromides, iodides, methane-, benzene- or p-toluenesulfonates. The etherification can be carried out, for example, in the presence of a base, an alkali metal hydride, hydroxide or carbonate, or of an amine. Conversely, corresponding ethers, such as lower alkoxy compounds, can be cleaved, for example, by means of strong acids, such as mineral acids, for example the hydrohalic acids hydrobromic or hydriodic acid, which may advantageously be present in the form of pyridinium halides, or by means of Lewis acids, for example halides of elements of main group III or the corresponding sub-groups. These reactions can be carried out, if necessary, with cooling or warming, for example in a temperature range from about -20° to about 100°C , in the presence or absence of a solvent or diluent, under inert gas and/or under pressure and, if appropriate, in a closed vessel.

Compounds according to the invention containing hydroxymethyl groups can be

prepared, for example, starting from compounds containing corresponding carboxyl or esterified carboxyl, corresponding compounds being reduced in a manner known per se, for example by reduction with a hydride which, if desired, may be complex, such as a hydride formed from an element of the 1st and 3rd main groups of the periodic table of the elements, for example borohydride or aluminohydride, for example lithium borohydride, lithium aluminium hydride, diisobutylaluminium hydride (an additional reduction step using alkali metal cyanoborohydride, such as sodium cyanoborohydride, may be necessary), and also diborane.

If an aromatic structural component is substituted by (lower) alkylthio (in $S(O)_n$ -R n is 0), this can be oxidised in a customary manner to corresponding (lower) alkanesulfinyl or -sulfonyl. Suitable oxidising agents for the oxidation to the sulfoxide step are, for example, inorganic peracids, such as peracids of mineral acids, for example periodic acid or persulfuric acid, organic peracids, such as appropriate percarboxylic or persulfonic acids, for example performic, peracetic, trifluoroperacetic or perbenzoic acid or p-toluenepersulfonic acid, or mixtures of hydrogen peroxide and acids, for example a mixture of hydrogen peroxide with acetic acid.

The oxidation is commonly carried out in the presence of suitable catalysts, catalysts which can be mentioned being suitable acids, such as substituted or unsubstituted carboxylic acids, for example acetic acid or trifluoroacetic acid, or transition metal oxides, such as oxides of elements of sub-group VII, for example vanadium oxide, molybdenum oxide or tungsten oxide. The oxidation is carried out under mild conditions, for example at temperatures from about -50° to about $+100^{\circ}\text{C}$.

The oxidation to the sulfone step may also be carried out appropriately at low temperatures using dinitrogen tetroxide as the catalyst in the presence of oxygen, just like the direct oxidation of (lower) alkylthio to (lower) alkanesulfonyl. However, in this case the oxidising agent is customarily employed in an excess.

If one of the variables contains amino, corresponding compounds of the formula (I), their tautomers or salts can be N-alkylated in a manner known per se; likewise, carbamoyl or radicals containing carbamoyl can be N-alkylated. The (aryl)alkylation is carried out, for example, using a reactive ester of an (aryl)C₁-C₇alkyl halide, for example a bromide or iodide, (aryl)C₁-C₇alkylsulfonate, for example methanesulfonate or p-toluenesulfonate, or a di-C₁-C₇alkyl sulfate, for example dimethyl sulfate, preferably under basic conditions, such as in the presence of sodium hydroxide solution or potassium hydroxide solution, and advantageously in the presence of a phase transfer catalyst, such as tetrabutylammonium bromide or benzyltrimethylammonium chloride, where, however, stronger basic condensing agents, such as alkali metal amides, hydrides or alkoxides, for example sodium amide, sodium hydride or sodium ethoxide, may be necessary. Amino can also be acylated in a manner known per se.

In compounds of the formula (I) which contain an esterified or amidated carboxyl group as a substituent, a group of this type can be converted into a free carboxyl group, for example by means of hydrolysis, for example in the presence of a basic agent, or of an acidic agent, such as a mineral acid. Tert-butyloxycarbonyl, for example, can furthermore be converted into carboxyl, for example in a manner known per se, such as treating with trihaloacetic acid, such as trifluoroacetic acid, and benzyloxycarbonyl can be converted into carboxyl, for example by catalytic hydrogenation in the presence of a hydrogenation catalyst, for example in the manner described below.

Furthermore, in compounds of the formula (I) which contain a carboxyl group as a substituent, this can be converted into an esterified carboxyl group, for example, by treating with an alcohol, such as a lower alkanol, in the presence of a suitable esterifying agent, such as an acid reagent, for example an inorganic or organic acid or a Lewis acid, for example zinc chloride, or a condensing agent which binds water, for example a carbodiimide, such as N,N'-dicyclohexylcarbodiimide, or by treating with a diazo reagent, such as with a diazo-lower alkane, for example diazomethane. This can also be obtained if compounds of the formula (I) in which the carboxyl group is present in free form or in salt form, such as

ammonium salt or metal salt form, for example alkali metal salt form, such as sodium salt or potassium salt form, are treated with a reactive ester of a (C₁-C₇)-alkyl halide, for example methyl or ethyl bromide or iodide, or an organic sulfonic acid ester, such as an appropriate (C₁-C₇)alkyl ester, for example methyl or ethyl methanesulfonate or p-toluenesulfonate.

Compounds of the formula (I) which contain an esterified carboxyl group as a substituent can be transesterified into other ester compounds of the formula (I) by transesterification, for example by treating with an alcohol, customarily a higher appropriate alcohol than that of the esterified carboxyl group in the starting material, in the presence of a suitable transesterifying agent, such as a basic agent, for example an alkali metal (C₁-C₇)alkanoate, (C₁-C₇)alkanolate or alkali metal cyanide, such as sodium acetate, sodium methoxide, sodium ethoxide, sodium tert-butoxide or sodium cyanide, or a suitable acid agent, if appropriate with removal of the resulting alcohol, for example by distillation. Appropriate, so-called activated esters of the formula (I) which contain an activated esterified carboxyl group as a substituent may also be used as starting materials (see below), and these may be converted into another ester by treating with a (C₁-C₇)-alkanol.

In compounds of the formula (I) which contain the carboxyl group as a substituent, this can also first be converted into a reactive derivative, such as an anhydride, including a mixed anhydride, such as an acid halide, for example an acid chloride (for example by treating with a thionyl halide, for example thionyl chloride), or an anhydride using a formic acid ester, for example a (C₁-C₇)alkyl ester (for example by treating a salt, such as an ammonium or alkali metal salt, with a haloformic acid ester, such as a chloroformic acid ester, such as a (C₁-C₇)alkyl ester), or into an activated ester, such as a cyanomethyl ester, a nitrophenyl ester, for example a 4-nitrophenyl ester, or a polyhalophenyl ester, for example a pentachlorophenyl ester (for example by treating with an appropriate hydroxyl compound in the presence of a suitable condensing agent, such as N,N'-dicyclohexylcarbodiimide), and then a reactive derivative of this type can be reacted with an amine and in this way amide compounds of the formula (I) which contain an amidated carboxyl group as a substituent can be obtained. In this

case, these can be obtained directly or via intermediate compounds; thus, for example, an activated ester, such as a 4-nitrophenyl ester, of a compound of the formula (I) containing a carboxyl group can first be reacted with a 1-unsubstituted imidazole and the 1-imidazolylcarbonyl compound obtained in this way brought to reaction with an amine. However, other non-activated esters, such as (C₁-C₇)alkyl esters of compounds of the formula (I), which contain, for example, (C₂-C₈)alkoxycarbonyl as a substituent, can also be brought to reaction with amines.

If an aromatic ring contains a hydrogen atom as a substituent, the latter can be replaced by a halogen atom with the aid of a halogenating agent in a customary manner, for example brominated with bromine, hypobromic acid, acyl hypobromites or other organic bromine compounds, for example N-bromosuccinimide, N-bromoacetamide, N-bromophthalimide, pyridinium perbromide, dioxane dibromide, 1,3-dibromo-5,5-dimethylhydantoin or 2,4,4,6-tetrabromo-2,5-cyclohexanedien-1-one, or chlorinated with elemental chlorine, for example in a halogenated hydrocarbon, such as chloroform, and with cooling, for example from down to about -10° to about +100°C.

If an aromatic ring in the compounds according to the invention contains an amino group, this can be diazotized in a customary manner, for example by treating with a nitrite, for example sodium nitrite, in the presence of a suitable protonic acid, for example a mineral acid, the reaction temperature advantageously being kept below about 5°C. The diazonium group present in the salt form and obtainable in this way can be substituted by analogous processes, for example as follows: by the hydroxyl group analogously to the boiling-out of phenol in the presence of water; by an alkoxy group by treating with an appropriate alcohol, energy having to be added; by the fluorine atom analogously to the Schiemann reaction in the thermolysis of corresponding diazonium tetrafluoroborates; by the halogen atoms chlorine, bromine or iodine and also the cyano group analogously to the Sandmeyer reaction in the reaction with corresponding Cu(I) salts, initially with cooling, for example to below about 5°C, and then heating, for example to about 60° to about 150°C.

If the compounds of the formula (I) contain unsaturated radicals, such as (lower) alkenyl or (lower) alkynyl groups, these can be converted into saturated radicals in a manner known per se. Thus, for example, multiple bonds are hydrogenated by catalytic hydrogenation in the presence of hydrogenation catalysts, suitable catalysts for this purpose being, for example, nickel, such as Raney nickel, and noble metals or their derivatives, for example oxides, such as palladium or platinum oxide, which may be applied, if desired, to support materials, for example to carbon or calcium carbonate. The hydrogenation may preferably be carried out at pressures between 1 and about 100 at and at room temperature between about -80° to about 200°C, in particular between room temperature and about 100°C. The reaction is advantageously carried out in a solvent, such as water, a lower alkanol, for example ethanol, isopropanol or n-butanol, an ether, for example dioxane, or a lower alkanecarboxylic acid, for example acetic acid.

Furthermore, in compounds of the formula (I) in which, for example, one of the aryl radicals contains halogen, such as chlorine, halogen can be replaced by reaction with a substituted or unsubstituted amine, an alcohol or a mercaptan.

The invention relates in particular to the processes described in the examples.

Salts of compounds of the formula (I) can be prepared in a manner known per se. Thus, for example, acid addition salts of compounds of the formula (I) are obtained by treating with an acid or a suitable ion exchange reagent. Salts can be converted into the free compounds in a customary manner, and acid addition salts can be converted, for example, by treating with a suitable basic agent.

Depending on the procedure and reaction conditions, the compounds according to the invention having salt-forming, in particular basic properties, can be obtained in free form or preferably in the form of salts.

In view of the close relationship between the novel compound in the free form and in the form of its salts, in the preceding text and below the free compound or its salts may correspondingly and advantageously also be understood as meaning the corresponding salts or the free compound.

The novel compounds including their salts of salt-forming compounds can also be obtained in the form of their hydrates or can include other solvents used for crystallization.

Depending on the choice of the starting materials and procedures, the novel compounds can be present in the form of one of the possible isomers or as mixtures thereof, for example as pure optical isomers, such as antipodes, or as isomer mixtures, such as racemates, diastereoisomer mixtures or racemate mixtures, depending on the number of asymmetric carbon atoms. For example, compounds of the formula (I) in which e.g. the heterocarbocyclic has an asymmetric C atom.

Racemates and diastereomer mixtures obtained can be separated into the pure isomers or racemates in a known manner on the basis of the physicochemical differences of the components, for example by fractional crystallization.

Racemates obtained may furthermore be resolved into the optical antipodes by known methods, for example by recrystallization from an optically active solvent, chromatography on chiral adsorbents, with the aid of suitable microorganisms, by cleavage with specific immobilized enzymes, via the formation of inclusion compounds, for example using chiral crown ethers, only one enantiomer being complexed, or by conversion into diastereomeric salts, for example by reaction of a basic final substance racemate with an optically active acid, such as a carboxylic acid, for example tartaric or malic acid, or sulfonic acid, for example camphorsulfonic acid, and separation of the diastereomer mixture obtained in this manner, for example on the basis of its differing solubilities, into the diastereomers from which the desired enantiomer can be liberated by the action of suitable agents. The more active enantiomer is advantageously isolated.

The invention also relates to those embodiments of the process, according to which a compound obtainable as an intermediate in any step of the process is used as a starting material and the missing steps are carried out or a starting material in the form of a derivative or salt and/or its racemates or antipodes is used or, in particular, formed under the reaction conditions.

In the process of the present invention, those starting materials are preferably used which lead to the compounds described as particularly useful at the beginning. The invention likewise relates to novel starting materials which have been specifically developed for the preparation of the compounds according to the invention, to their use and to processes for their preparation, the variables alk, R₁, R₂, R₃, and X having the meanings indicated for the preferred compound groups of the formula (I) in each case.

The invention likewise relates to pharmaceutical preparations which contain the compounds according to the invention or pharmaceutically acceptable salts thereof as active ingredients, and to processes for their preparation.

The pharmaceutical preparations according to the invention which contain the compound according to the invention or pharmaceutically acceptable salts thereof are those for enteral, such as oral, furthermore rectal, and parenteral administration to (a) warm-blooded animal(s), the pharmacological active ingredient being present on its own or together with a pharmaceutically acceptable carrier. The daily dose of the active ingredient depends on the age and the individual condition and also on the manner of administration.

The novel pharmaceutical preparations contain, for example, from about 10 % to about 80%, preferably from about 20 % to about 60 %, of the active ingredient. Pharmaceutical preparations according to the invention for enteral or parenteral administration are, for example, those in unit dose forms, such as sugar-coated tablets, tablets, capsules or suppositories, and furthermore ampoules. These are prepared in a manner known per se, for example by means of conventional mixing, granulating, sugar-coating, dissolving or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by combining the active ingredient with solid carriers, if desired granulating a mixture obtained, and processing the mixture or granules, if desired or necessary, after addition of suitable excipients to give tablets or sugar-coated tablet cores.

Suitable carriers are, in particular, fillers, such as sugars, for example lactose, sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, furthermore binders, such as starch paste, using, for example, corn, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose and/or polyvinylpyrrolidone, if desired, disintegrants, such as the abovementioned starches, furthermore carboxymethyl starch, crosslinked polyvinylpyrrolidone, agar, alginic acid or a salt thereof, such as sodium alginate; auxiliaries are primarily glidants, flow-regulators and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol. Sugar-coated tablet cores are provided with suitable coatings which, if desired, are resistant to gastric juice, using, inter alia, concentrated sugar solutions which, if desired, contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, coating solutions in suitable organic solvents or solvent mixtures or, for the preparation of gastric juice-resistant coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Colorants or pigments, for example to identify or to indicate different doses of active ingredient, may be added to the tablets or sugar-coated tablet coatings.

Other orally utilizable pharmaceutical preparations are hard gelatin capsules, and also soft closed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The hard gelatin capsules may contain the active ingredient in the form of granules, for example in a mixture with fillers, such as lactose, binders, such as starches, and/or lubricants, such as talc or magnesium stearate, and, if desired, stabilizers. In soft capsules, the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols, it also being possible to add stabilizers.

Suitable rectally utilizable pharmaceutical preparations are, for example, suppositories, which consist of a combination of the active ingredient with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols. Furthermore, gelatin rectal capsules which contain a combination of the

active ingredient with a base substance may also be used. Suitable base substances are, for example, liquid triglycerides, polyethylene glycols or paraffin hydrocarbons.

Suitable preparations for parenteral administration are primarily aqueous solutions of an active ingredient in water-soluble form, for example a water-soluble salt, and furthermore suspensions of the active ingredient, such as appropriate oily injection suspensions, using suitable lipophilic solvents or vehicles, such as fatty oils, for example sesame oil, or synthetic fatty acid esters, for example ethyl oleate or triglycerides, or aqueous injection suspensions which contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if necessary, also stabilizers.

The dose of the active ingredient depends on the warm-blooded animal species, the age and the individual condition and on the manner of administration. In the normal case, an approximate daily dose of about 10 mg to about 250 mg is to be estimated in the case of oral administration for a patient weighing approximately 75 kg .

The following examples illustrate the invention described above; however, they are not intended to limit its extent in any manner. Temperatures are indicated in degrees Celsius.

The following examples illustrate the invention.

Abbreviations:

HCl	hydrochloric acid
NaOH	sodium hydroxide
THF	tetrahydrofuran
min	minute(s)
h	hour(s)
m.p.	melting point


FAB-MS Fast Atom Bombardment Mass Spectroscopy

Rf retention factor on a thin layer chromatography plate

Solvent systems (v/v/v):

A1:	hexanes / ethyl acetate	1:1
A2:	hexanes / ethyl acetate	2:1
B1:	dichloromethane / methanol	9:1
C1:	dichloromethane / methanol / ammonium hydroxide	90:10:1

Example 1: Naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-yl]-ethyl}-amide hydrochloride



A suspension of 2-chloro-quinazolin-4-ylamine (0.282 g) and naphthalene-1-sulfonic acid (2-piperidin-4-yl-ethyl)-amide (0.5 g) in isopentylalcohol (10 ml) is heated up to 120 °C for 15 h. The resulting solution is concentrated and chromatographed (silica gel, B1) to give product as a foam. This material is taken up in dichloromethane (100 ml) and treated at 0 °C with a 4 N HCl solution in dioxane (2 ml). Concentration *in vacuo*, followed by crystallization from ethyl acetate and methanol gives naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-yl]-ethyl}-amide hydrochloride melting at 184 - 191 °C. Rf(B1) 0.30; FAB-MS: (M+H)⁺ = 462.

The starting material can be prepared, for example, as follows:

a) 4-[2-(Naphthalene-1-sulfonylamino)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester


A solution of 4-(2-amino-ethyl)-piperidine-1-carboxylic acid *tert*-butyl ester (1.5 g) and diisopropylethylamine (1.69 ml) in acetonitrile (30 ml) is cooled to 0 °C and treated with a

solution of naphthalene-1-sulfonylchloride (1.64 g) in acetonitrile (10 ml). The reaction is stirred at ambient temperature for 24 h and concentrated *in vacuo*. The residue is taken up in ethyl acetate, washed with a 0.5 N HCl solution, a saturated aqueous sodium carbonate solution and water. It is then dried over sodium sulfate and concentrated to a tan powder. Chromatography on silica gel (A2) affords 4-[2-(naphthalene-1-sulfonylamino)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester, melting at 50 - 53 °C. Rf(A1) 0.40.

b) Naphthalene-1-sulfonic acid (2-piperidin-4-yl-ethyl)-amide

A suspension of 4-[2-(naphthalene-1-sulfonylamino)-ethyl]-piperidine-1-carboxylic acid *tert*-butyl ester (2.25 g) in dichloromethane (10 ml) is treated with a 4 N HCl solution in dioxane (10 ml) at 0 °C. Under completion, the reaction mixture is concentrated *in vacuo*, the residue is taken up in a 1 N sodium hydroxide solution and dichloromethane. After extraction with dichloromethane, the organics are dried over sodium sulfate and concentrated to give naphthalene-1-sulfonic acid (2-piperidin-4-yl-ethyl)-amide. Rf(C3) 0.12.

Example 2: Naphthalene-1-sulfonic acid [1-(4-amino-quinazolin-2-yl)-piperidin-4-ylmethyl]-amide hydrochloride



Following the procedure described in Example 1, a mixture of 0.236 g of 2-chloro-quinazolin-4-ylamine and 0.4 g of naphthalene-1-sulfonic acid (piperidin-4-yl-methyl)-amide is converted to naphthalene-1-sulfonic acid [1-(4-amino-quinazolin-2-yl)-piperidin-4-ylmethyl]-amide hydrochloride, melting at 268 - 272 °C. Rf(B1) 0.46; FAB-MS: (M+H)⁺ = 448.


The starting material can be prepared, for example, as follows:

a) 4-[(Naphthalene-1-sulfonylamino)-methyl]-piperidine-1-carboxylic acid *tert*-butyl ester

Following the procedure described in Example 1a, a mixture of 1.89 g of 4-(amino-methyl-piperidine-1-carboxylic acid *tert*-butyl ester (obtained from 4-aminomethyl-piperidine as described in: *Synth. Commun.* **1992**, 22, 2357) and 2 g of naphthalene-1-sulfonylchloride in acetonitrile gives 4-[(naphthalene-1-sulfonylamino)-methyl]-piperidine-1-carboxylic acid *tert*-butyl ester as a foam. Rf(A2) 0.20.

b) Naphthalene-1-sulfonic acid (piperidin-4-yl-methyl)-amide

Following the procedure described in Example 1b, 3.12 g of 4-[(naphthalene-1-sulfonylamino)-methyl]-piperidine-1-carboxylic acid *tert*-butyl ester are converted to naphthalene-1-sulfonic acid (piperidin-4-yl-methyl)-amide, melting at 141 - 143. Rf(C1) 0.15.

Example 3: Naphthalene-1-sulfonic acid {2-[4-(4-amino-quinazolin-2-yl)-piperazin-1-yl]-ethyl}-amide hydrochloride

Following the procedure described in Example 1, a mixture of 0.282 g of 2-chloro-quinazolin-4-ylamine and 0.5 g of naphthalene-1-sulfonic acid (2-piperidin-4-yl-ethyl)-amide is converted to naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-yl]-ethyl}-amide hydrochloride, melting at 184 - 191 °C. Rf(B2) 0.30; FAB-MS: (M+H)⁺ = 462.

The starting material can be prepared, for example, as follows:


a) 4-[2-(Naphthalene-1-sulfonylamino)-ethyl]-piperazine-1-carboxylic acid *tert*-butyl ester

Following the procedure described in Example 1a, a mixture of 1.82 g of 4-(2-amino-ethyl)-piperazine-1-carboxylic acid *tert*-butyl ester (obtained from 1-(2-amino-ethyl)-piperazine as described in: *Synth. Commun.* **1992**, 22, 2357) and 1.8 g of naphthalene-1-sulfonylchloride in acetonitrile gives 4-[2-(naphthalene-1-sulfonylamino)-ethyl]-piperazine-1-carboxylic acid *tert*-butyl ester as a foam. Rf(A1) 0.33.

b) Naphthalene-1-sulfonic acid (2-piperazin-1-yl-ethyl)-amide

Following the procedure described in Example 1, 2.86 g of 4-[2-(naphthalene-1-sulfonylamino)-ethyl]-piperazine-1-carboxylic acid *tert*-butyl ester are converted to naphthalene-1-sulfonic acid (2-piperazin-1-yl-ethyl)-amide, melting at 152 - 154 °C. Rf(C2) 0.41.

Example 4: 2-[4-(Naphthalene-1-sulfonyl)-piperazin-1-yl]-quinazolin-4-ylamine hydrochloride



Following the procedure described in Example 1, a mixture of 0.3 g of 2-chloro-quinazolin-4-ylamine and 0.462 g of 1-(naphthalene-1-sulfonyl)-piperazine is converted to 2-[4-(naphthalene-1-sulfonyl)-piperazin-1-yl]-quinazolin-4-ylamine hydrochloride, melting at 254 - 264 °C. Rf(C4) 0.68; FAB-MS: (M+H)⁺ = 420.

The starting material can be prepared, for example, as follows:

a) 4-(Naphthalene-1-sulfonyl)-piperazine-1-carboxylic acid *tert*-butyl ester

Following the procedure described in Example 1a, a mixture of 1.56 g of piperazine-N-carbamic acid *tert*-butyl ester and 1.9 g of naphthalene-1-sulfonylchloride in acetonitrile

- 64 -

gives 4-(naphthalene-1-sulfonyl)-piperazine-1-carboxylic acid *tert*-butyl ester, melting at 143 - 145 °C. Rf(A5) 0.27.

b) 1-(Naphthalene-1-sulfonyl)-piperazine

Following the procedure described in Example 1b, 2.91 g of 4-(naphthalene-1-sulfonyl)-piperazine-1-carboxylic acid *tert*-butyl ester are converted to 1-(naphthalene-1-sulfonyl)-piperazine, melting at 85 - 87 °C. Rf(C1) 0.55.

Example 5: Tablets, each containing 50 mg of active ingredient, for example, naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-yl]-ethyl}-amide hydrochloride, can be prepared as follows:

Composition (for 10,000 tablets)

Active ingredient	500.0 g
Lactose	500.0 g
Potato starch	352.0 g
Gelatin	8.0 g
Talc	60.0 g
Magnesium stearate	10.0 g
Silica (highly disperse)	20.0 g
Ethanol	q.s.

The active ingredient is mixed with the lactose and 292 g of potato starch, and the mixture is moistened using an alcoholic solution of the gelatin and granulated by means of a sieve. After drying, the remainder of the potato starch, the talc, the magnesium stearate and the highly disperse silica are admixed and the mixture is compressed to give tablets of weight 145.0 mg each and active ingredient content 50.0 mg which, if desired, can be provided with breaking notches for finer adjustment of the dose.

Example 6: Coated tablets, each containing 100 mg of active ingredient, for example, naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-

yl]-ethyl]-amide hydrochloride, can be prepared as follows:

Composition (for 1000 tablets):

Active ingredient	100.00 g
Lactose	100.00 g
Corn starch	70.00 g
Talc	8.50 g
Calcium stearate	1.50 g
Hydroxypropylmethylcellulose	2.36 g
Shellac	0.64 g
Water	q.s.
Dichloromethane	q.s.

The active ingredient, the lactose and 40 g of the corn starch are mixed and moistened and granulated with a paste prepared from 15 g of corn starch and water (with warming). The granules are dried, and the remainder of the corn starch, the talc and the calcium stearate are added and mixed with the granules. The mixture is compressed to give tablets (weight: 280 mg) and these are coated with a solution of the hydroxypropylmethylcellulose and the shellac in dichloromethane (final weight of the coated tablet: 283 mg).

Example 7: Tablets and coated tablets containing another compound of the formula (I) or a pharmaceutically acceptable salt of a compound of the formula (I), for example as in one of Examples 1 to 4, can also be prepared in an analogous manner to that described in Examples 5 and 6.

SEQUENCE LISTING

(1) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1501 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 61..1432

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```
TTAGTTTTGT TCTGAGAACG TTAGAGTTAT AGTACCGTGC GATCGTTCTT CAAGCTGCTA      60
ATG GAC GTC CTC TTC TTC CAC CAG GAT TCT AGT ATG GAG TTT AAG CTT      108
Met Asp Val Leu Phe Phe His Gln Asp Ser Ser Met Glu Phe Lys Leu
  1             5             10             15
GAG GAG CAT TTT AAC AAG ACA TTT GTC ACA GAG AAC AAT ACA GCT GCT      156
Glu Glu His Phe Asn Lys Thr Phe Val Thr Glu Asn Asn Thr Ala Ala
```

- 67 -

	20		25		30	
GCT CGG AAT GCA GCC TTC CCT GCC TGG GAG GAC TAC AGA GGC AGC GTA						204
Ala Arg Asn Ala Ala Phe Pro Ala Trp Glu Asp Tyr Arg Gly Ser Val						
	35		40		45	
GAC GAT TTA CAA TAC TTT CTG ATT GGG CTC TAT ACA TTC GTA AGT CTT						252
Asp Asp Leu Gln Tyr Phe Leu Ile Gly Leu Tyr Thr Phe Val Ser Leu						
	50		55		60	
CTT GGC TTT ATG GGC AAT CTA CTT ATT TTA ATG GCT GTT ATG AAA AAG						300
Leu Gly Phe Met Gly Asn Leu Leu Ile Leu Met Ala Val Met Lys Lys						
	65		70		75	80
CGC AAT CAG AAG ACT ACA GTG AAC TTT CTC ATA GGC AAC CTG GCC TTC						348
Arg Asn Gln Lys Thr Thr Val Asn Phe Leu Ile Gly Asn Leu Ala Phe						
		85		90		95
TCC GAC ATC TTG GTC GTC CTG TTT TGC TCC CCT TTC ACC CTG ACC TCT						396
Ser Asp Ile Leu Val Val Leu Phe Cys Ser Pro Phe Thr Leu Thr Ser						
	100		105		110	
GTC TTG TTG GAT CAG TGG ATG TTT GGC AAA GCC ATG TGC CAT ATC ATG						444
Val Leu Leu Asp Gln Trp Met Phe Gly Lys Ala Met Cys His Ile Met						
	115		120		125	
CCG TTC CTT CAA TGT GTG TCA GTT CTG GTT TCA ACT CTG ATT TTA ATA						492
Pro Phe Leu Gln Cys Val Ser Val Leu Val Ser Thr Leu Ile Leu Ile						
	130		135		140	
TCA ATT GCC ATT GTC AGG TAT CAT ATG ATA AAG CAC CCT ATT TCT AAC						540
Ser Ile Ala Ile Val Arg Tyr His Met Ile Lys His Pro Ile Ser Asn						
	145		150		155	160
AAT TTA ACG GCA AAC CAT GGC TAC TTC CTG ATA GCT ACT GTC TGG ACA						588
Asn Leu Thr Ala Asn His Gly Tyr Phe Leu Ile Ala Thr Val Trp Thr						
		165		170		175
CTG GGC TTT GCC ATC TGT TCT CCC CTC CCA GTG TTT CAC AGT CTT GTG						636
Leu Gly Phe Ala Ile Cys Ser Pro Leu Pro Val Phe His Ser Leu Val						
	180		185		190	
GAA CTT AAG GAG ACC TTT GGC TCA GCA CTG CTG AGT AGC AAA TAT CTC						684

- 68 -

Glu	Leu	Lys	Glu	Thr	Phe	Gly	Ser	Ala	Leu	Leu	Ser	Ser	Lys	Tyr	Leu		
	195						200					205					
TGT	GTT	GAG	TCA	TGG	CCC	TCT	GAT	TCA	TAC	AGA	ATT	GCT	TTC	ACA	ATC	732	
Cys	Val	Glu	Ser	Trp	Pro	Ser	Asp	Ser	Tyr	Arg	Ile	Ala	Phe	Thr	Ile		
	210					215					220						
TCT	TTA	TTG	CTA	GTG	CAG	TAT	ATC	CTG	CCT	CTA	GTA	TGT	TTA	ACG	GTA	780	
Ser	Leu	Leu	Leu	Val	Gln	Tyr	Ile	Leu	Pro	Leu	Val	Cys	Leu	Thr	Val		
225					230					235					240		
AGT	CAT	ACC	AGC	GTC	TGC	CGA	AGC	ATA	AGC	TGT	GGA	TTG	TCC	CAC	AAA	828	
Ser	His	Thr	Ser	Val	Cys	Arg	Ser	Ile	Ser	Cys	Gly	Leu	Ser	His	Lys		
				245					250					255			
GAA	AAC	AGA	CTC	GAA	GAA	AAT	GAG	ATG	ATC	AAC	TTA	ACC	CTA	CAG	CCA	876	
Glu	Asn	Arg	Leu	Glu	Glu	Asn	Glu	Met	Ile	Asn	Leu	Thr	Leu	Gln	Pro		
			260					265					270				
TCC	AAA	AAG	AGC	AGG	AAC	CAG	GCA	AAA	ACC	CCC	AGC	ACT	CAA	AAG	TGG	924	
Ser	Lys	Lys	Ser	Arg	Asn	Gln	Ala	Lys	Thr	Pro	Ser	Thr	Gln	Lys	Trp		
		275					280						285				
AGC	TAC	TCA	TTC	ATC	AGA	AAG	CAC	AGA	AGG	AGG	TAC	AGC	AAG	AAG	ACG	972	
Ser	Tyr	Ser	Phe	Ile	Arg	Lys	His	Arg	Arg	Arg	Tyr	Ser	Lys	Lys	Thr		
	290					295					300						
GCC	TGT	GTC	TTA	CCC	GCC	CCA	GCA	GGA	CCT	TCC	CAG	GGG	AAG	CAC	CTA	1020	
Ala	Cys	Val	Leu	Pro	Ala	Pro	Ala	Gly	Pro	Ser	Gln	Gly	Lys	His	Leu		
305					310					315					320		
GCC	GTT	CCA	GAA	AAT	CCA	GCC	TCC	GTC	CGT	AGC	CAG	CTG	TCG	CCA	TCC	1068	
Ala	Val	Pro	Glu	Asn	Pro	Ala	Ser	Val	Arg	Ser	Gln	Leu	Ser	Pro	Ser		
				325					330					335			
AGT	AAG	GTC	ATT	CCA	GGG	GTC	CCA	ATC	TGC	TTT	GAG	GTG	AAA	CCT	GAA	1116	
Ser	Lys	Val	Ile	Pro	Gly	Val	Pro	Ile	Cys	Phe	Glu	Val	Lys	Pro	Glu		
			340					345					350				
GAA	AGC	TCA	GAT	GCT	CAT	GAG	ATG	AGA	GTC	AAG	CGT	TCC	ATC	ACT	AGA	1164	
Glu	Ser	Ser	Asp	Ala	His	Glu	Met	Arg	Val	Lys	Arg	Ser	Ile	Thr	Arg		
		355					360					365					

- 69 -

ATA AAA AAG AGA TCT CGA AGT GTT TTC TAC AGA CTG ACC ATA CTG ATA	1212
Ile Lys Lys Arg Ser Arg Ser Val Phe Tyr Arg Leu Thr Ile Leu Ile	
370 375 380	
CTC GTG TTC GCC GTT AGC TGG ATG CCA CTC CAC GTC TTC CAC GTG GTG	1260
Leu Val Phe Ala Val Ser Trp Met Pro Leu His Val Phe His Val Val	
385 390 395 400	
ACT GAC TTC AAT GAT AAC TTG ATT TCC AAT AGG CAT TTC AAG CTG GTA	1308
Thr Asp Phe Asn Asp Asn Leu Ile Ser Asn Arg His Phe Lys Leu Val	
405 410 415	
TAC TGC ATC TGT CAC TTG TTA GGC ATG ATG TCC TGT TGT CTA AAT CCG	1356
Tyr Cys Ile Cys His Leu Leu Gly Met Met Ser Cys Cys Leu Asn Pro	
420 425 430	
ATC CTA TAT GGT TTC CTT AAT AAT GGT ATC AAA GCA GAC TTG AGA GCC	1404
Ile Leu Tyr Gly Phe Leu Asn Asn Gly Ile Lys Ala Asp Leu Arg Ala	
435 440 445	
CTT ATC CAC TGC CTA CAC ATG TCA TGA TTCTCTCTGTG CACCAAAGAG	1452
Leu Ile His Cys Leu His Met Ser *	
450 455	
AGAAGAAACG TGGTAATTGA CACATAATTT ATACAGAAGT ATTCTGGAT	1501

- 70 -

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 457 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Asp Val Leu Phe Phe His Gln Asp Ser Ser Met Glu Phe Lys Leu
 1             5             10             15

Glu Glu His Phe Asn Lys Thr Phe Val Thr Glu Asn Asn Thr Ala Ala
          20             25             30

Ala Arg Asn Ala Ala Phe Pro Ala Trp Glu Asp Tyr Arg Gly Ser Val
          35             40             45

Asp Asp Leu Gln Tyr Phe Leu Ile Gly Leu Tyr Thr Phe Val Ser Leu
 50             55             60

Leu Gly Phe Met Gly Asn Leu Leu Ile Leu Met Ala Val Met Lys Lys
 65             70             75             80

Arg Asn Gln Lys Thr Thr Val Asn Phe Leu Ile Gly Asn Leu Ala Phe
          85             90             95

Ser Asp Ile Leu Val Val Leu Phe Cys Ser Pro Phe Thr Leu Thr Ser
          100            105            110

Val Leu Leu Asp Gln Trp Met Phe Gly Lys Ala Met Cys His Ile Met
          115            120            125

Pro Phe Leu Gln Cys Val Ser Val Leu Val Ser Thr Leu Ile Leu Ile
          130            135            140

Ser Ile Ala Ile Val Arg Tyr His Met Ile Lys His Pro Ile Ser Asn
          145            150            155            160

```


- 71 -

Asn	Leu	Thr	Ala	Asn	His	Gly	Tyr	Phe	Leu	Ile	Ala	Thr	Val	Trp	Thr	165	170	175
Leu	Gly	Phe	Ala	Ile	Cys	Ser	Pro	Leu	Pro	Val	Phe	His	Ser	Leu	Val	180	185	190
Glu	Leu	Lys	Glu	Thr	Phe	Gly	Ser	Ala	Leu	Leu	Ser	Ser	Lys	Tyr	Leu	195	200	205
Cys	Val	Glu	Ser	Trp	Pro	Ser	Asp	Ser	Tyr	Arg	Ile	Ala	Phe	Thr	Ile	210	215	220
Ser	Leu	Leu	Leu	Val	Gln	Tyr	Ile	Leu	Pro	Leu	Val	Cys	Leu	Thr	Val	225	230	235
Ser	His	Thr	Ser	Val	Cys	Arg	Ser	Ile	Ser	Cys	Gly	Leu	Ser	His	Lys	245	250	255
Glu	Asn	Arg	Leu	Glu	Glu	Asn	Glu	Met	Ile	Asn	Leu	Thr	Leu	Gln	Pro	260	265	270
Ser	Lys	Lys	Ser	Arg	Asn	Gln	Ala	Lys	Thr	Pro	Ser	Thr	Gln	Lys	Trp	275	280	285
Ser	Tyr	Ser	Phe	Ile	Arg	Lys	His	Arg	Arg	Arg	Tyr	Ser	Lys	Lys	Thr	290	295	300
Ala	Cys	Val	Leu	Pro	Ala	Pro	Ala	Gly	Pro	Ser	Gln	Gly	Lys	His	Leu	305	310	315
Ala	Val	Pro	Glu	Asn	Pro	Ala	Ser	Val	Arg	Ser	Gln	Leu	Ser	Pro	Ser	325	330	335
Ser	Lys	Val	Ile	Pro	Gly	Val	Pro	Ile	Cys	Phe	Glu	Val	Lys	Pro	Glu	340	345	350
Glu	Ser	Ser	Asp	Ala	His	Glu	Met	Arg	Val	Lys	Arg	Ser	Ile	Thr	Arg	355	360	365
Ile	Lys	Lys	Arg	Ser	Arg	Ser	Val	Phe	Tyr	Arg	Leu	Thr	Ile	Leu	Ile	370	375	380

- 72 -

Leu Val Phe Ala Val Ser Trp Met Pro Leu His Val Phe His Val Val
385 390 395 400

Thr Asp Phe Asn Asp Asn Leu Ile Ser Asn Arg His Phe Lys Leu Val
405 410 415

Tyr Cys Ile Cys His Leu Leu Gly Met Met Ser Cys Cys Leu Asn Pro
420 425 430

Ile Leu Tyr Gly Phe Leu Asn Asn Gly Ile Lys Ala Asp Leu Arg Ala
435 440 445

Leu Ile His Cys Leu His Met Ser *
450 455

(3) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1457 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 61..1432

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

- 73 -

GTTTCCCTCT GAATAGATTA ATTTAAAGTA GTCATGTAAT GTTTTTTTGG TTGCTGACAA	60
ATG TCT TTT TAT TCC AAG CAG GAC TAT AAT ATG GAT TTA GAG CTC GAC	108
Met Ser Phe Tyr Ser Lys Gln Asp Tyr Asn Met Asp Leu Glu Leu Asp	
1 5 10 15	
GAG TAT TAT AAC AAG ACA CTT GCC ACA GAG AAT AAT ACT GCT GCC ACT	156
Glu Tyr Tyr Asn Lys Thr Leu Ala Thr Glu Asn Asn Thr Ala Ala Thr	
20 25 30	
CGG AAT TCT GAT TTC CCA GTC TGG GAT GAC TAT AAA AGC AGT GTA GAT	204
Arg Asn Ser Asp Phe Pro Val Trp Asp Asp Tyr Lys Ser Ser Val Asp	
35 40 45	
GAC TTA CAG TAT TTT CTG ATT GGG CTC TAT ACA TTT GTA AGT CTT CTT	252
Asp Leu Gln Tyr Phe Leu Ile Gly Leu Tyr Thr Phe Val Ser Leu Leu	
50 55 60	
GGC TTT ATG GGG AAT CTA CTT ATT TTA ATG GCT CTC ATG AAA AAG CGT	300
Gly Phe Met Gly Asn Leu Leu Ile Leu Met Ala Leu Met Lys Lys Arg	
65 70 75 80	
AAT CAG AAG ACT ACG GTA AAC TTC CTC ATA GGC AAT CTG GCC TTT TCT	348
Asn Gln Lys Thr Thr Val Asn Phe Leu Ile Gly Asn Leu Ala Phe Ser	
85 90 95	
GAT ATC TTG GTT GTG CTG TTT TGC TCA CCT TTC ACA CTG ACG TCT GTC	396
Asp Ile Leu Val Val Leu Phe Cys Ser Pro Phe Thr Leu Thr Ser Val	
100 105 110	
TTG CTG GAT CAG TGG ATG TTT GGC AAA GTC ATG TGC CAT ATT ATG CCT	444
Leu Leu Asp Gln Trp Met Phe Gly Lys Val Met Cys His Ile Met Pro	
115 120 125	
TTT CTT CAA TGT GTG TCA GTT TTG GTT TCA ACT TTA ATT TTA ATA TCA	492
Phe Leu Gln Cys Val Ser Val Leu Val Ser Thr Leu Ile Leu Ile Ser	
130 135 140	
ATT GCC ATT GTC AGG TAT CAT ATG ATA AAA CAT CCC ATA TCT AAT AAT	540
Ile Ala Ile Val Arg Tyr His Met Ile Lys His Pro Ile Ser Asn Asn	
145 150 155 160	

- 74 -

TTA ACA GCA AAC CAT GGC TAC TTT CTG ATA GCT ACT GTC TGG ACA CTA	588
Leu Thr Ala Asn His Gly Tyr Phe Leu Ile Ala Thr Val Trp Thr Leu	
165 170 175	
GGT TTT GCC ATC TGT TCT CCC CTT CCA GTG TTT CAC AGT CTT GTG GAA	636
Gly Phe Ala Ile Cys Ser Pro Leu Pro Val Phe His Ser Leu Val Glu	
180 185 190	
CTT CAA GAA ACA TTT GGT TCA GCA TTG CTG AGC AGC AGG TAT TTA TGT	684
Leu Gln Glu Thr Phe Gly Ser Ala Leu Leu Ser Ser Arg Tyr Leu Cys	
195 200 205	
GTT GAG TCA TGG CCA TCT GAT TCA TAC AGA ATT GCC TTT ACT ATC TCT	732
Val Glu Ser Trp Pro Ser Asp Ser Tyr Arg Ile Ala Phe Thr Ile Ser	
210 215 220	
TTA TTG CTA GTT CAG TAT ATT CTG CCC TTA GTT TGT CTT ACT GTA AGT	780
Leu Leu Leu Val Gln Tyr Ile Leu Pro Leu Val Cys Leu Thr Val Ser	
225 230 235 240	
CAT ACA AGT GTC TGC AGA AGT ATA AGC TGT GGA TTG TCC AAC AAA GAA	828
His Thr Ser Val Cys Arg Ser Ile Ser Cys Gly Leu Ser Asn Lys Glu	
245 250 255	
AAC AGA CTT GAA GAA AAT GAG ATG ATC AAC TTA ACT CTT CAT CCA TCC	876
Asn Arg Leu Glu Glu Asn Glu Met Ile Asn Leu Thr Leu His Pro Ser	
260 265 270	
AAA AAG AGT GGG CCT CAG GTG AAA CTC TCT GGC AGC CAT AAA TGG AGT	924
Lys Lys Ser Gly Pro Gln Val Lys Leu Ser Gly Ser His Lys Trp Ser	
275 280 285	
TAT TCA TTC ATC AAA AAA CAC AGA AGA AGA TAT AGC AAG AAG ACA GCA	972
Tyr Ser Phe Ile Lys Lys His Arg Arg Arg Tyr Ser Lys Lys Thr Ala	
290 295 300	
TGT GTG TTA CCT GCT CCA GAA AGA CCT TCT CAA GAG AAC CAC TCC AGA	1020
Cys Val Leu Pro Ala Pro Glu Arg Pro Ser Gln Glu Asn His Ser Arg	
305 310 315 320	
ATA CTT CCA GAA AAC TTT GGC TCT GTA AGA AGT CAG CTC TCT TCA TCC	1068
Ile Leu Pro Glu Asn Phe Gly Ser Val Arg Ser Gln Leu Ser Ser Ser	
325 330 335	

- 75 -

AGT AAG TTC ATA CCA GGG GTC CCC ACT TGC TTT GAG ATA AAA CCT GAA	1116
Ser Lys Phe Ile Pro Gly Val Pro Thr Cys Phe Glu Ile Lys Pro Glu	
340 345 350	
GAA AAT TCA GAT GTT CAT GAA TTG AGA GTA AAA CGT TCT GTT ACA AGA	1164
Glu Asn Ser Asp Val His Glu Leu Arg Val Lys Arg Ser Val Thr Arg	
355 360 365	
ATA AAA AAG AGA TCT CGA AGT GTT TTC TAC AGA CTG ACC ATA CTG ATA	1212
Ile Lys Lys Arg Ser Arg Ser Val Phe Tyr Arg Leu Thr Ile Leu Ile	
370 375 380	
TTA GTA TTT GCT GTT AGT TGG ATG CCA CTA CAC CTT TTC CAT GTG GTA	1260
Leu Val Phe Ala Val Ser Trp Met Pro Leu His Leu Phe His Val Val	
385 390 395 400	
ACT GAT TTT AAT GAC AAT CTT ATT TCA AAT AGG CAT TTC AAG TTG GTG	1308
Thr Asp Phe Asn Asp Asn Leu Ile Ser Asn Arg His Phe Lys Leu Val	
405 410 415	
TAT TGC ATT TGT CAT TTG TTG GGC ATG ATG TCC TGT TGT CTT AAT CCA	1356
Tyr Cys Ile Cys His Leu Leu Gly Met Met Ser Cys Cys Leu Asn Pro	
420 425 430	
ATT CTA TAT GGG TTT CTT AAT AAT GGG ATT AAA GCT GAT TTA GTG TCC	1404
Ile Leu Tyr Gly Phe Leu Asn Asn Gly Ile Lys Ala Asp Leu Val Ser	
435 440 445	
CTT ATA CAC TGT CTT CAT ATG TAA TAA TTCTCACTGT TTACCAAGGA	1452
Leu Ile His Cys Leu His Met * *	
450 455	
AAGAAC	1457

(4) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 457 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

- 76 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Ser	Phe	Tyr	Ser	Lys	Gln	Asp	Tyr	Asn	Met	Asp	Leu	Glu	Leu	Asp	1	5	10	15
Glu	Tyr	Tyr	Asn	Lys	Thr	Leu	Ala	Thr	Glu	Asn	Asn	Thr	Ala	Ala	Thr	20	25	30	
Arg	Asn	Ser	Asp	Phe	Pro	Val	Trp	Asp	Asp	Tyr	Lys	Ser	Ser	Val	Asp	35	40	45	
Asp	Leu	Gln	Tyr	Phe	Leu	Ile	Gly	Leu	Tyr	Thr	Phe	Val	Ser	Leu	Leu	50	55	60	
Gly	Phe	Met	Gly	Asn	Leu	Leu	Ile	Leu	Met	Ala	Leu	Met	Lys	Lys	Arg	65	70	75	80
Asn	Gln	Lys	Thr	Thr	Val	Asn	Phe	Leu	Ile	Gly	Asn	Leu	Ala	Phe	Ser	85	90	95	
Asp	Ile	Leu	Val	Val	Leu	Phe	Cys	Ser	Pro	Phe	Thr	Leu	Thr	Ser	Val	100	105	110	
Leu	Leu	Asp	Gln	Trp	Met	Phe	Gly	Lys	Val	Met	Cys	His	Ile	Met	Pro	115	120	125	
Phe	Leu	Gln	Cys	Val	Ser	Val	Leu	Val	Ser	Thr	Leu	Ile	Leu	Ile	Ser	130	135	140	
Ile	Ala	Ile	Val	Arg	Tyr	His	Met	Ile	Lys	His	Pro	Ile	Ser	Asn	Asn	145	150	155	160
Leu	Thr	Ala	Asn	His	Gly	Tyr	Phe	Leu	Ile	Ala	Thr	Val	Trp	Thr	Leu	165	170	175	
Gly	Phe	Ala	Ile	Cys	Ser	Pro	Leu	Pro	Val	Phe	His	Ser	Leu	Val	Glu	180	185	190	

- 77 -

Leu	Gln	Glu	Thr	Phe	Gly	Ser	Ala	Leu	Leu	Ser	Ser	Arg	Tyr	Leu	Cys	195	200	205	
Val	Glu	Ser	Trp	Pro	Ser	Asp	Ser	Tyr	Arg	Ile	Ala	Phe	Thr	Ile	Ser	210	215	220	
Leu	Leu	Leu	Val	Gln	Tyr	Ile	Leu	Pro	Leu	Val	Cys	Leu	Thr	Val	Ser	225	230	235	240
His	Thr	Ser	Val	Cys	Arg	Ser	Ile	Ser	Cys	Gly	Leu	Ser	Asn	Lys	Glu	245	250	255	
Asn	Arg	Leu	Glu	Glu	Asn	Glu	Met	Ile	Asn	Leu	Thr	Leu	His	Pro	Ser	260	265	270	
Lys	Lys	Ser	Gly	Pro	Gln	Val	Lys	Leu	Ser	Gly	Ser	His	Lys	Trp	Ser	275	280	285	
Tyr	Ser	Phe	Ile	Lys	Lys	His	Arg	Arg	Arg	Tyr	Ser	Lys	Lys	Thr	Ala	290	295	300	
Cys	Val	Leu	Pro	Ala	Pro	Glu	Arg	Pro	Ser	Gln	Glu	Asn	His	Ser	Arg	305	310	315	320
Ile	Leu	Pro	Glu	Asn	Phe	Gly	Ser	Val	Arg	Ser	Gln	Leu	Ser	Ser	Ser	325	330	335	
Ser	Lys	Phe	Ile	Pro	Gly	Val	Pro	Thr	Cys	Phe	Glu	Ile	Lys	Pro	Glu	340	345	350	
Glu	Asn	Ser	Asp	Val	His	Glu	Leu	Arg	Val	Lys	Arg	Ser	Val	Thr	Arg	355	360	365	
Ile	Lys	Lys	Arg	Ser	Arg	Ser	Val	Phe	Tyr	Arg	Leu	Thr	Ile	Leu	Ile	370	375	380	
Leu	Val	Phe	Ala	Val	Ser	Trp	Met	Pro	Leu	His	Leu	Phe	His	Val	Val	385	390	395	400
Thr	Asp	Phe	Asn	Asp	Asn	Leu	Ile	Ser	Asn	Arg	His	Phe	Lys	Leu	Val	405	410	415	

- 78 -


Tyr Cys Ile Cys His Leu Leu Gly Met Met Ser Cys Cys Leu Asn Pro
420 425 430

Ile Leu Tyr Gly Phe Leu Asn Asn Gly Ile Lys Ala Asp Leu Val Ser
435 440 445

Leu Ile His Cys Leu His Met * *
450

What is claimed is

1. Use of a compound of formula (I),



in which

alk represents a single bond or lower alkylene;

the integer n is 0 or 1;

R₁ represents

- (i) hydrogen, halogen, nitro, cyano, lower alkyl, lower alkenyl, lower alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, by lower alkoxy, by amino, by substituted amino, by carboxy, by lower alkoxycarbonyl, by (carbocyclic or heterocyclic) aryl-lower alkoxycarbonyl, by carbamoyl, or by N-substituted carbamoyl;
- (ii) amino or substituted amino;
- (iii) hydroxy, lower alkoxy, lower alkenyloxy, lower alkynyloxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, C₃-C₈-cycloalkoxy, C₃-C₈-cycloalkyl-lower alkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, lower alkoxycarbonyl-oxy, (carbocyclic or heterocyclic) aryl-lower alkoxycarbonyl-oxy, aminocarbonyl-oxy, or N-substituted aminocarbonyl-oxy;
- (iv) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (v) carbamoyl or N-substituted carbamoyl;
- (vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-O-R, -NR₀₁-CO-R, -NR₀₁-CO-NR₀₁-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R and R₀₁ being as defined below, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O, S(O)_n or NR₀ or which may be condensed at two adjacent carbon atoms with a benzene ring}]; or

(vii) an element of formula $-X_1(X_2)(X_3)$ wherein, (a) if X_1 is $-\text{CH}-$, X_2 together with X_3 represent a structural element of formula $-X_4(\text{CO})_p(\text{CH}_2)_o-$, $-(\text{CH}_2)_q-X_4(\text{CO})_p(\text{CH}_2)_r-$, or $-(\text{CH}_2)_s-X_4\text{CO}-(\text{CH}_2)_t-$; or, (b) if X_1 is $-\text{N}-$, X_2 together with X_3 represent a structural element of formula $-\text{CO}-(\text{CH}_2)_u-$; [X_4 being $-\text{CH}_2-$, $-\text{N}(\text{R}_{01})-$ or $-\text{O}-$; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X_4 is different from $-\text{CH}_2-$];

R_2 and R_3 , independently of one another, represent

(i) hydrogen, lower alkyl, lower alkenyl, lower alkynyl, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
 (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, N-substituted carbamoyl, and $-\text{S}(\text{O})_n\text{-R}$;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O , $\text{S}(\text{O})_n$, NR_0 or which may be condensed at two adjacent carbon atoms with a benzene ring];

X represents N or CH ;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, lower alkenyl, lower alkynyl, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, lower alkynyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, (carbocyclic or heterocyclic) aroyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl-lower alkyl}$, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy,

lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iv) amino, substituted amino;

(v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, S(O)_n or NR₀ or which may be condensed at two adjacent carbon atoms with a benzene ring], or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen, lower alkyl, lower alkenyl, lower alkinyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower alkanoyl, (carbocyclic or heterocyclic) aroyl, -SO₂-R, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;

wherein in each case R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy; or a pharmaceutically acceptable salt thereof for the manufacture of a pharmaceutical composition for the prophylaxis and treatment of diseases or disorders associated with NPY Y5 receptor subtype.

2. Use according to claim 1 for the manufacture of a pharmaceutical composition for the prophylaxis and the treatment of disorders or disease states caused by eating disorders, of obesity, bulimia nervosa, diabetes, dyslipidemia, hypertension, memory loss, epileptic seizures, migraine, sleep disturbance, pain, sexual/reproductive disorders, depression, anxiety, cerebral hemorrhage, shock, congestive heart failure, nasal congestion or diarrhea.

3. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

- (i) hydrogen, halogen, nitro, cyano, lower alkyl, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by lower alkoxy, by substituted amino, by lower alkoxy-carbonyl, or by N-substituted carbamoyl;
- (ii) substituted amino;
- (iii) hydroxy, lower alkoxy, lower alkoxy-lower alkoxy, C₃-C₈-cycloalkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, lower alkoxy-carbonyl-oxy, or N-substituted aminocarbonyl-oxy;
- (iv) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (v) N-substituted carbamoyl;
- (vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-O-R, -NR₀₁-CO-R, -NR₀₁-CO-NR₁-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R being as defined below and R₀₁ being as defined above, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O, S(O)_n or NR₀} or which is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring]; or
- (vii) an element of formula -X₁(X₂)(X₃) wherein, (a) if X₁ is -CH-, X₂ together with X₃ represent a structural element of formula -X₄-(CO)_p-(CH₂)_o-, -(CH₂)_q-X₄-(CO)_p-(CH₂)_r-, or -(CH₂)_s-X₄-CO-(CH₂)_t-; or, (b) if X₁ is -N-, X₂ together with X₃ represent a structural element of formula -CO-(CH₂)_u-; [X₄ being -CH₂-, -N(R₀₁)- or -O-; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X₄ is different from -CH₂-];

R₂ and R₃, independently of one another, represent

- (i) hydrogen, lower alkyl, lower alkenyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, N-substituted carbamoyl, and -S(O)_n-R;

R₂ and R₃ together represent lower alkylene [which may be interrupted by O, S(O)_n, or NR₀] or represent lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring;

X represents N or CH; and the integer n is 0 or 1;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

(i) halogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, nitro, cyano;

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryloxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;

(iv) amino, substituted amino;

(v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, is derived and selected from the group consisting of phenyl, biphenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furyl, thienyl, pyridyl, indolyl, indazolyl, benzofuryl, benzothiophenyl, benzimidazolyl, quinoliny, isoquinoliny, or quinazoliny;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O,

S(O)_n or NR₀] or is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring, or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen or lower alkyl;

wherein, in each case, R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;.

4. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

(i) hydrogen, halogen, nitro, cyano, lower alkyl, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryl, or lower alkyl which is substituted by halogen, by lower alkoxy, by substituted amino, by lower alkoxy-carbonyl, or by substituted carbamoyl;

(ii) substituted amino;

(iii) hydroxy, lower alkoxy, C₃-C₈-cycloalkoxy, (carbocyclic or heterocyclic) aryl-lower alkoxy, or lower alkoxy-lower alkoxy;

(iv) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, or (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;

(v) N-substituted carbamoyl;

(vi) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, -SO₂-NR₀₁-R, or -SO₂-NR₀₁-CO-R, [R and R₁ being as defined below, or the group -N(R)(R₀₁) represents amino which is di-substituted by lower alkylene {which may be interrupted by O or NR₀} or which is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring]; or

(vii) an element of formula -X₁(X₂)(X₃) wherein, (a) if X₁ is -CH-, X₂ together with X₃ represent a structural element of formula -X₄-(CO)_p-(CH₂)_o-, -(CH₂)_q-X₄-(CO)_p-(CH₂)_r-, or -(CH₂)_s-X₄-CO-(CH₂)_t-; or, (b) if X₁ is -N-, X₂ together with X₃ represent a structural element of formula -CO-(CH₂)_u-; [X₄ being -CH₂-, -N(R₀₁)- or -O-; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X₄ is different from -CH₂-];

R_2 and R_3 , independently of one another, represent

- (i) hydrogen, lower alkyl, lower alkenyl, (carbocyclic or heterocyclic) aryl, or (carbocyclic or heterocyclic) aryl-lower alkyl;
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: lower alkoxy, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, and substituted carbamoyl;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O, S(O)_n, or NR₀];

X represents N or CH; and the integer n is 0 or 1;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

- (i) halogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, lower alkoxy, lower alkenyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, nitro, cyano;
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, (carbocyclic or heterocyclic) aryloxy, amino, substituted amino, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iv) substituted amino;
- (v) lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, is derived from phenyl, naphthyl or pyridyl;

wherein, in each case, the substituted amino group of substituted amino, of N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-

- 86 -

cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, S(O)_n or NR₀] or is di-substituted by lower alkylene which is condensed at two adjacent carbon atoms with a benzene ring, or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen or lower alkyl;

wherein, in each case, R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy.

5. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents

- (i) hydrogen, halogen, cyano, nitro, lower alkyl, C₃-C₈-cycloalkyl, or phenyl;
- (ii) amino which is mono-substituted by lower alkyl, phenyl or pyridyl, or which is disubstituted by lower alkyl or by C₂-C₆-alkylene;
- (iii) hydroxy or lower alkoxy which is unsubstituted or substituted by C₃-C₈-cycloalkyl or by phenyl;
- (iv) a group selected from -CH(OH)-R, -CO-R, -NR₀₁-CO-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, or -SO₂-NR₀₁-R, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or naphthyl, R₀₁ being hydrogen or lower alkyl, or the group -N(R)(R₀₁) represents amino which is mono-substituted by lower alkyl, by hydroxy-lower alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C₂-C₆-alkylene {which may be interrupted by O or NR₀, R₀ being hydrogen or lower alkyl}];

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

R₃ represents hydrogen;

X represents N or CH; and the integer n is 0 or 1;

wherein any aryl moiety, if not designated otherwise, and the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting

of halogen, lower alkyl, halo-alkyl, lower alkoxy, hydroxy, hydroxy-lower alkoxy, and lower alkoxy-lower alkoxy;.

6. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents a single bond or lower alkylene;

R₁ represents (i) hydrogen, halogen, cyano, nitro, lower alkyl, C₃-C₈-cycloalkyl, or phenyl;

(ii) amino which is mono-substituted by lower alkyl, phenyl or pyridyl, or which is disubstituted by lower alkyl or by C₂-C₆-alkylene;

(iii) hydroxy or lower alkoxy which is unsubstituted or substituted by C₃-C₈-cycloalkyl, or by phenyl;

(iv) a group selected from -NR₀₁-CO-R, -NR₀₁-SO₂-R, -NR₀₁-SO₂-NR₀₁-R, -SO₂-R, or -SO₂-NR₀₁-R, [R being lower alkyl, halo-lower alkyl, phenyl, pyridyl, or naphthyl, R₀₁ being hydrogen or lower alkyl, or the group -N(R)(R₀₁) represents amino which is mono-substituted by lower alkyl, by hydroxy-lower alkyl, or by naphthyl, or which is di-substituted by lower alkyl or by C₂-C₆-alkylene {which may be interrupted by O or NR₀, R₀ being hydrogen or lower alkyl}];

R₂ represents hydrogen or phenyl which is unsubstituted or is substituted by a substituent selected from the group consisting of: halogen, cyano, lower alkyl, lower alkoxy, and oxy-lower alkylene-oxy;

R₃ represents hydrogen;

X represents N or CH; and the integer n is 0 or 1;

wherein any aryl moiety, if not designated otherwise, and the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of halogen, lower alkyl, halo-lower alkyl, lower alkoxy, hydroxy, and hydroxy-lower alkoxy;.

7. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents -(CH₂)_n- and the integer n is 1-3;

R₁ represents the group of formula NH-SO₂-R and R is naphthyl;

R₂ and R₃ each are hydrogen;

X is N or CH; and

the ring A is unsubstituted or substituted by C₁-C₄-alkoxy.

8. Use according to claim 1 or 2 of a compound of formula (I) or a pharmaceutically acceptable salt thereof in which

alk represents $-(CH_2)_o-$ and the integer o is 1 or 2;

R_1 represents the group of formula $NH-SO_2-R$ and R is naphthyl;

R_2 and R_3 each are hydrogen;

X is CH; and

the ring A is unsubstituted.

9. A compound of formula (I) or a salt thereof in which;

alk represents a single bond or lower alkylene;

the integer n is 0 or 1;

R_1 represents

(vi) a group selected from $-CH(OH)-R$, $-CO-R$, $-NR_{01}-CO-O-R$, $-NR_{01}-CO-R$, $-NR_{01}-CO-NR_{01}-R$, $-NR_{01}-SO_2-R$, $-NR_{01}-SO_2-NR_{01}-R$, $-SO_2-R$, $-SO_2-NR_{01}-R$, or $-SO_2-NR_{01}-CO-R$, [R and R_1 being as defined below, or the group $-N(R)(R_{01})$ represents amino which is di-substituted by lower alkylene {which may be interrupted by O, $S(O)_n$ or NR_0 or which may be condensed at two adjacent carbon atoms with a benzene ring}]; or

(vii) an element of formula $-X_1(X_2)(X_3)$ wherein, (a) if X_1 is $-CH-$, X_2 together with X_3 represent a structural element of formula $-X_4-(CO)_p-(CH_2)_o-$, $-(CH_2)_q-X_4-(CO)_p-(CH_2)_r-$, or $-(CH_2)_s-X_4-CO-(CH_2)_t-$; or, (b) if X_1 is $-N-$, X_2 together with X_3 represent a structural element of formula $-CO-(CH_2)_u-$; [X_4 being $-CH_2-$, $-N(R_{01})-$ or $-O-$; the integer o is 3-5; the integer p is 0 or 1; the integer q is 1 or 2; the integer r is 1; the integer s is 1 or 2; the integer t is 1 or 2; the integer u is 3-5; with the proviso that, if the integer p is 0, X_4 is different from $-CH_2-$];

R_2 and R_3 , independently of one another, represent

(i) hydrogen, lower alkyl, lower alkenyl, lower alkynyl, C_3-C_8 -cycloalkyl, C_3-C_8 -cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl; or

(ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, hydroxy-lower alkoxy, lower alkoxy-lower alkoxy, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, N-substituted carbamoyl, and $-S(O)_n-R$;

R_2 and R_3 together represent lower alkylene [which may be interrupted by O, $S(O)_n$, NR_0 or which may be condensed at two adjacent carbon atoms with a benzene ring];

X represents N or CH;

wherein, in each case, any aryl moiety, for example, of (carbocyclic or heterocyclic) aryl, aroyl, or aryloxy, respectively, as well as the benzo ring A is unsubstituted or substituted by one or more substituents selected from the group consisting of

- (i) halogen, lower alkyl, lower alkenyl, lower alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, lower alkoxy, lower alkenyloxy, lower alkynyloxy, oxy-lower alkylene-oxy, hydroxy, lower alkanoyloxy, (carbocyclic or heterocyclic) aryl-lower alkanoyloxy, lower alkanoyl, (carbocyclic or heterocyclic) aryl-lower alkanoyl, (carbocyclic or heterocyclic) aroyl, nitro, cyano;
- (ii) lower alkyl which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iii) lower alkoxy which is substituted by a substituent selected from the group consisting of: halogen, hydroxy, lower alkoxy, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryloxy, (carbocyclic or heterocyclic) aryl, amino, substituted amino, carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl, carbamoyl, and N-substituted carbamoyl;
- (iv) amino, substituted amino;
- (v) carboxy, lower alkoxy-carbonyl, lower alkoxy-lower alkoxy-carbonyl, (carbocyclic or heterocyclic) aryl-lower alkoxy-carbonyl;
- (vi) carbamoyl and N-substituted carbamoyl;

wherein, in each case, the substituted amino group of substituted amino, N-substituted carbamoyl, and of N-substituted aminocarbonyl-oxy is (i) mono-substituted or, independently of one another, di-substituted by lower alkyl, by C₃-C₈-cycloalkyl, by C₃-C₈-cycloalkyl-lower alkyl, by (carbocyclic or heterocyclic) aryl, by (carbocyclic or heterocyclic) aryl-lower alkyl, or is (ii) di-substituted by lower alkylene [which may be interrupted by O, S(O)_n or NR₀ or which may be condensed at two adjacent carbon atoms with a benzene ring], or is (iii) mono-substituted or, in the second line, independently of one another, di-substituted by -CO-(O)_v-R and the integer v is 0 or 1;

wherein, in each case, the integer n is 0, 1 or 2;

wherein, in each case, R₀ represents hydrogen, lower alkyl, lower alkenyl, lower alkynyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, lower

- 90 -

alkanoyl, (carbocyclic or heterocyclic) aroyl, -SO₂-R, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;

wherein, in each case, R₀₁ represents hydrogen or lower alkyl;

wherein, in each case, R represents hydrogen, lower alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-lower alkyl, (carbocyclic or heterocyclic) aryl, (carbocyclic or heterocyclic) aryl-lower alkyl, or lower alkyl which is substituted by halogen, by hydroxy, or by lower alkoxy;

10. A compound according to claim 9 of formula (I) or a salt thereof selected from the group consisting of

Naphthalene-1-sulfonic acid {2-[1-(4-amino-quinazolin-2-yl)-piperidin-4-yl]-ethyl}-amide;

Naphthalene-1-sulfonic acid [1-(4-amino-quinazolin-2-yl)-piperidin-4-ylmethyl]-amide;

Naphthalene-1-sulfonic acid {2-[4-(4-amino-quinazolin-2-yl)-piperazin-1-yl]-ethyl}-amide; and

2-[4-(Naphthalene-1-sulfonyl)-piperazin-1-yl]-quinazolin-4-ylamine.

11. A pharmaceutical composition for the treatment of diseases or disorders associated with NPY Y5 receptor subtype comprising a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof according to claim 1 and a carrier.

12. A method for the treatment and prophylaxis of disorders or disease states associated with NPY Y5 receptor subtype comprising administering to a warm-blooded animal, including man, in need of such treatment a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof according to claim 1.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 96/05055

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D239/95 C07D401/04 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 511 836 A (H-J. HESS ET AL.) 12 May 1970 see the whole document ---	1-9
X	US 3 635 979 A (H-J. HESS ET AL.) 18 January 1972 see the whole document ---	1-9
X	FR 2 321 890 A (SYNTHELABO) 25 March 1977 see page 1 - page 15; table 1 -----	1-9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

8 April 1997

Date of mailing of the international search report

16.04.97

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 96/05055

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 12
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 96/05055

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3511836 A	12-05-70	NONE	
US 3635979 A	18-01-72	US 3663706 A	16-05-72
FR 2321890 A	25-03-77	NONE	